


SPEAKER NOTES

MODULE 05 – TACTICAL TRAUMA ASSESSMENT

SLIDE 1 – TITLE SLIDE

This module is an overview of tactical trauma assessment (TTA). The skills practice will take place near the end of the course after you have learned ALL of the skills.



SLIDE 2 – CHANGE LOG

The Curriculum Change Log serves as a centralized reference to quickly track recent updates to training materials. It supports trainers by promoting clear communication, accountability, and alignment, helping stakeholders and learners understand what changes were made, why they were implemented, and when they occurred.

TACTICAL TRAUMA ASSESSMENT

CHANGE LOG - Curriculum Update History

CHANGE DATE	PRODUCT UPDATE	DESCRIPTION OF CHANGE
29 April 2025	Module 05 - Didactic PPT Slides 26-27, and 29	Updated TCCC 1380 Card image to reflect changing "gender" to "sex"

TCCC-CLS-PPT-02_29 APR 25 2

SLIDE 3 – TCCC ROLES

Tactical Combat Casualty Care is broken up into four roles of care. The most basic is taught to All Service Members (ASM), which is designed to instruct in the absolute basics of hemorrhage control and to recognize more serious injuries.

You are in the Combat Lifesaver (CLS) role. This teaches you more advanced care to treat the most common causes of death on the battlefield, and to recognize, prevent, and communicate with medical personnel the life-threatening complications of these injuries.

The Combat Medic/Corpsman (CMC) role includes much more advanced and invasive care requiring significantly more medical knowledge and skills.

Finally, the last role, Combat Paramedic/Provider (CPP) is for Combat paramedics and advanced providers, to provide the most sophisticated care to keep our wounded warriors alive and get them to definitive care.

Your role as a CLS is to treat the most common causes of death on the battlefield, which are massive hemorrhage and airway/respiratory problems. Also, you are given the skills to prevent complications and treat other associated but not immediately life-threatening injuries.



TCCC-CLS-PPT-02_29 APR 25 2

SPEAKER NOTES

SLIDE 4 – TLO/ELO

The Tactical Trauma Assessment (TTA) module has three cognitive learning objectives and five performance learning objectives. The cognitive learning objectives are to identify the:

1. Common causes of altered mental status in combat or noncombat environments
2. Importance of disarming and securing communications equipment of a casualty with altered mental status
3. Importance and techniques of communicating with a casualty in TFC

The performance learning objectives are to demonstrate:

1. Techniques used to assess a casualty for responsiveness
2. Communication with a casualty in Tactical Field Care
3. Application of body substance isolation (BSI) in TFC
4. TTA in the proper order using the MARCH PAWS sequence in accordance with CoTCCC guidelines
5. Appropriate actions and interventions used during a casualty assessment to render aid to the casualty in accordance with CoTCCC Guidelines.

The critical aspects are to identify the importance of and demonstrate the systematic approach for assessment and interventions in providing lifesaving care to a casualty following the MARCH PAWS sequence in accordance with the CoTCCC guidelines.

SLIDE 5 – MARCH PAWS

A full tactical trauma assessment should follow the MARCH PAWS sequence.

- **M**assive bleeding
- **A**irway
- **R**espiration/breathing
- **C**irculation
- **H**ypothermia/Head injuries
- **P**ain
- **A**ntibiotics
- **W**ounds
- **S**plinting

We will cover the interventions and procedures of MARCH PAWS in more detail in later modules.

CLS
TCCC

STUDENT LEARNING OBJECTIVES

LEARNING OBJECTIVES

06 Given a combat or noncombat scenario, perform a Tactical Trauma Assessment (TTA) in accordance with CoTCCC Guidelines.

- 29 Demonstrate the techniques used to assess a casualty for responsiveness
- 30 Identify the common causes of altered mental status in combat or noncombat environments
- 31 Identify the importance of disarming and securing communications equipment of a casualty with altered mental status
- 32 Identify the importance and techniques of communicating with a casualty in Tactical Field Care
- 33 Demonstrate communicating with a casualty in Tactical Field Care
- 34 Demonstrate application of body substance isolation (BSI) in Tactical Field Care
- 35 Demonstrate a Tactical Trauma Assessment in the proper order using the MARCH PAWS sequence in accordance with CoTCCC guidelines
- 36 Demonstrate the appropriate actions and interventions used during a casualty assessment to render aid to the casualty in accordance with CoTCCC Guidelines

01 **TERMINAL LEARNING OBJECTIVES (TLOs)** 08 **ENABLING LEARNING OBJECTIVES (ELOs)**

● = Performance ELOs
● = Cognitive ELOs

TCCC CLS-PPT-06-29 APR 21 4

CLS
TCCC

TACTICAL FIELD CARE

MARCH PAWS

DURING LIFE-THREATENING	AFTER LIFE-THREATENING
M MASSIVE BLEEDING #1 Priority	P PAIN
A AIRWAY	A ANTIBIOTICS
R RESPIRATION (breathing)	W WOUNDS
C CIRCULATION	S SPLINTING
H HYPOTHERMIA / HEAD INJURIES	


TCCC CLS-PPT-06-29 APR 21 5

SPEAKER NOTES

SLIDE 6 – TACTICAL TRAUMA ASSESSMENT HOW-TO (VIDEO)


Play video

Pay attention to this video. You will be expected to perform a full TTA upon completion of this training.




SLIDE 7 – COMBAT SPEED TTA "FIRE FIGHT CONSCIOUS CASUALTY" (VIDEO)

Play video



SLIDE 8 – COMBAT SPEED TTA "EXPLOSION" UNCONSCIOUS CASUALTY (VIDEO)

Play video



SLIDE 9 – BODY SUBSTANCE ISOLATION (BSI)

Whenever possible, the responder/CLS should don latex-free gloves as a BSI precaution.

Gloves are provided in the JFAK and CLS bags.

TACTICAL TRAUMA ASSESSMENT BODY SUBSTANCE ISOLATION (BSI)



Whenever possible, the responder should don **latex-free** gloves as a precaution

TCCC-CLS-PPT-09_29 APR 25 9

SPEAKER NOTES

SLIDE 10 – CASUALTY BLOOD SWEEP

Your initial casualty evaluation should be a rapid head-to-toe check for any unrecognized life-threatening bleeding (a blood sweep).

This blood sweep should include a visual and hands-on (palpation) inspection of the front and back of the casualty from head to toe, including neck, armpits, groin, etc.

SLIDE 11 – QUICKLY IDENTIFY MASSIVE, LIFE-THREATENING BLEEDING

This blood sweep is a systematic way to ensure rapid identification of any unrecognized life-threatening bleeding.

SLIDE 12 – HEMORRHAGE CONTROL





If you identify life-threatening bleeding that was missed in the Care Under Fire phase, immediately apply a tourniquet or hemostatic dressing, and/or pressure dressing.

If a tourniquet was previously applied but bleeding is not controlled, apply a second tourniquet side-by-side with the original tourniquet, preferably higher on the injured limb, if possible, to control the bleeding. This is the “M” of MARCH PAWS.

SLIDE 13 – IDENTIFYING OBSTRUCTED AIRWAY

Evaluate the casualty’s airway and ensure the airway is open. **LOOK** (for rise and fall of the chest), **LISTEN** (for sounds of breathing), and **FEEL** (breath on your cheek) for indications of trouble breathing, snoring or gurgling sounds, visible objects obstructing the airway, and any severe trauma to the face.

Do not do a blind finger sweep.



SPEAKER NOTES

SLIDE 14 – IN A CASUALTY WITHOUT AN AIRWAY OBSTRUCTION, YOU CAN PERFORM THE FOLLOWING MANEUVERS TO OPEN THE AIRWAY

If a casualty is unconscious, the tongue may have relaxed, causing an airway blockage. Use the head-tilt chin-lift or jaw-thrust method to open the airway.

Important note: If a neck or spinal injury is suspected, use the jaw-thrust method to open the airway.

This is the first “A” of MARCH PAWS.

NOTE: Once the airway has been opened using one of these maneuvers, the casualty may require repeated/continued maneuvers to maintain an open airway.



SLIDE 15 – MANAGEMENT/RECOVERY POSITION

If the casualty is **conscious**, allow them to assume any position that best protects the airway and allows them to breath easily, including sitting up.

Place an **unconscious casualty** in the recovery position. If an NPA was inserted into the right nostril, place the casualty on their right side, if possible.



SLIDE 16 – RESPIRATIONS


LOOK (for rise and fall of the chest), **LISTEN** (for sounds of breathing), and **FEEL** (breath on your cheek) for indications of trouble breathing (as noted for airway previously).

Respiration rate (breaths per minute) and quality (shallow, labored, etc.) should be noted.

Indications of respiratory distress include:

- Breathing that is progressively difficult
- Decreased breathing sounds
- Distended neck veins
- Opposed to “progressive” respiratory distress
- Hunched over; they need to be in the “position of comfort”
- Agitation due to a lack of oxygen
- High pulse

This is the “R” of MARCH PAWS.



SPEAKER NOTES

SLIDE 17 – LIFE-THREATENING CHEST INJURY


Common causes of chest injuries include gunshot, stab, or shrapnel wounds to chest and blunt-force trauma.

Note obvious signs of penetrating trauma, bruising, swelling, crackling/popping (on palpation), or other deformities of the chest. **Check** the casualty's respiration and ability to breathe. All open and/or sucking chest wounds should be treated by immediately applying a vented chest seal to cover the defect.

For respiratory distress not resolved by a chest seal or in a casualty with known or suspected chest or back trauma without an open and/or sucking chest wound, consider a tension pneumothorax, and perform a needle decompression of the chest.

Injuries to the chest are very serious and can be life-threatening. The casualty's condition can change quickly with a chest injury.

RESPIRATION ASSESSMENT AND MANAGEMENT IN TFC LIFE-THREATENING CHEST INJURY




TCCC-CLS-PPT-01_29 APR 23 17

SLIDE 18 – REASSESS TREATMENTS

This is the “C” in MARCH PAWS. The casualty should be reassessed for life-threatening hemorrhage (including effectiveness of prior interventions-TQs, pressure bandages, etc.).

Is there an obvious pelvic or femur fracture? If so, a medic should be informed immediately. Assess the radial pulse.

If the pulse is absent or weak, shock should be suspected, and a medic should be informed immediately.




TCCC-CLS-PPT-02_29 APR 23 18

SLIDE 19 – GENERAL INDICATOR OF SHOCK

The CLS should be familiar with the signs/symptoms of shock. In the combat environment, shock is assumed to be due to blood loss.

If untreated, shock could lead to death. If shock is suspected, a medic should be informed immediately.




TCCC-CLS-PPT-03 29 APR 23 19

SLIDE 20 – HYPOTHERMIA PREVENTION

Prevent hypothermia by minimizing the casualty's exposure to the elements and applying active hypothermia prevention measures, when possible.

If no rewarming equipment is available, then use dry blankets, poncho liners, sleeping bags, or anything that will retain heat and keep the casualty dry. Make sure you assess for hemorrhagic shock and ensure bleeding is controlled.




TCCC-CLS-PPT-04 29 APR 23 20

SPEAKER NOTES

SLIDE 21 – IF A PENETRATING EYE INJURY IS NOTED OR SUSPECTED

Perform rapid field test of visual acuity (e.g., read Meals Ready to Eat (MRE) label, or name tag).

If the casualty has any penetrating injuries, they should take the antibiotic in the Combat Wound Medication Pack (CWMP). Cover eye with rigid shield, not pressure patch. **Do not** cover both eyes unless both are injured and you are sure the casualty will not return to the fight.



SLIDE 22 – COMBAT WOUND MEDICATION PACK

The CWMP contains drugs for mild to moderate pain (meloxicam and acetaminophen) and an antibiotic specific for penetrating wounds (moxifloxacin).

A CWMP can give significant pain relief for mild to moderate pain and will not alter the casualty's mental status. It also includes antibiotics for preventing/treating infections after traumatic injuries, such as penetrating wounds, eye injuries, and burns.




This is the “P” and “A” of MARCH PAWS.

SLIDE 23 – INSPECT AND ADDRESS ALL KNOWN WOUNDS

This is the “W” in MARCH PAWS.

All other wounds (burns, fractures, other soft tissue wounds, etc.) should be addressed with splinting, dressings, etc. as appropriate. This will be covered in more detail in a later module.

Note: Reassess pulses after all dressings are placed to ensure that they are not too tight. Do not ever apply one and forget it!




SLIDE 24 – BURN CARE

Stop the burning process by extracting the casualty from the source, and cover the burned areas with dry, sterile dressings.

If the burn is caused by **white phosphorus**, submerge the affected area in water, if possible; otherwise, the dressing must be wet. Advise medical personnel immediately.

Remember to prioritize assessing MARCH before addressing burns. This is part of the “W” of MARCH PAWS.



SPEAKER NOTES

SLIDE 25 – ASSESS FOR A FRACTURE

Assess for **any fractures**, and if present, splint the fracture using whatever materials are available, making sure to immobilize the joint above and the joint below the fracture.

Check pulse(s) before and after applying splints. Treat open fractures with meds (for pain and to prevent infection) with meds from the CWMP.

This is the “**S**” of MARCH PAWS.

SLIDE 26 – COMMUNICATION AND DOCUMENTATION

Communicate with the casualty by reassuring them and telling them about procedures being performed.

Communicate with medical personnel and your **tactical leadership**, and relay casualty status and evacuation needs.

SLIDE 27 – PHASE 3: TACTICAL EVACUATION CARE

Document all assessment and care on a DD Form 1380 TCCC card. Every Service member will carry their own DD Form 1380 in their JFAK. If possible, use a permanent marker (such as a Sharpie) to make entries on the card. When rendering care or assisting medical personnel, include as much information as you can on the card.

This is the official record of the care provided and should go with the casualty when care is handed off to a medic or at the time of evacuation.

Communication includes the **MIST report** and **MEDEVAC request**.


SLIDE 28 – SKILL STATION

At this time, we will break for a trainer-led demonstration on TTA.

CLS
TCCC

SECONDARY INJURIES

ASSESS FOR A FRACTURE



CLOSED FRACTURE OPEN FRACTURE

WARNING SIGNS OF A FRACTURE:


- Significant pain and swelling
- An audible or perceived “snap”
- Different length or shape of limb
- Loss of pulse or sensation in the injured arm or leg (check pulse before and after treatment)
- Crepitus (hearing a crackling or popping sound under the skin)

TCCC-CLS-PPT-06_29 APR 25 25

CLS
TCCC

TACTICAL FIELD CARE

COMMUNICATION



Communicate with the casualty if possible

- Encourage.
- Reassure.
- Explain care each step of the way

Communicate with tactical leadership as soon as possible with status and evacuation requirements throughout casualty treatment as needed

COMMUNICATE WITH EVACUATION AND MEDICAL ASSETS

- Communicate with the evacuation system to coordinate TACEVAC/MEDEVAC using the MEDEVAC request
- Keep the casualty's DD Form 1380 up to date

TACTICAL COMBAT CASUALTY CARE (TCCC) CARD

BATTLE NUMBER: _____

NAME: _____ LAST NAME: _____

DOB: (MM/DD) _____ SEX: (M/F) _____

UNIT: _____ ALLIED: _____

TCCC-CLS-PPT-06_29 APR 25 26

CLS
TCCC

TACTICAL EVACUATION

PHASE 3: TACTICAL EVACUATION CARE

CASUALTY MONITORING

- Continue to reassess and monitor casualty

EVAC REQUEST


- Uses 9-Line format

CASUALTY PREP

- Secure items
- Prep litter
- Prep evac equipment
- Pack casualty

COMPLETE MIST REPORT


- M Mechanism of injury
- I Injuries
- S Symptoms
- T Treatment



(Litter Transport)

PRE-EVAC PROCEDURES

- Complete DD Form 1380



TCCC-CLS-PPT-06_29 APR 25 27

CLS
TCCC

TRAINER-LED DEMONSTRATION


- Tactical Trauma Assessment

TCCC-CLS-PPT-06_29 APR 25 28

SPEAKER NOTES

SLIDE 29 – SUMMARY

In this module, we discussed the TTA. We identified the common causes of altered mental status in combat or noncombat environments, the importance of disarming and securing the communications equipment of a casualty with altered mental status, and techniques for communicating with a casualty in TFC. We also demonstrated techniques for assessing a casualty for responsiveness, applying body substance isolation, conducting a TTA in the proper order using the MARCH PAWS sequence, and using appropriate actions and interventions in a casualty assessment to render aid in accordance with CoTCCC Guidelines.




SLIDE 30 – CHECK ON LEARNING

Ask questions of the learners referring to key concepts from the module.

Now for a check on learning:

1. In which phase of care is the TTA performed?
 - TFC
2. What pneumonic is used to prioritize care in the TTA?
 - MARCH PAWS
3. What is a blood sweep?
 - A blood sweep is your initial casualty evaluation. It should be a rapid head-to-toe check for any unrecognized life-threatening bleeding.



SLIDE 31 – QUESTIONS

