



U.S. CENTRAL COMMAND COVID-19 PANDEMIC PLAYBOOK FOR OPERATIONAL ENVIRONMENTS



This playbook contains guidance and references generally available as of the issuance date and is meant to be a reference for pandemic response. It does not supersede the Department of Defense Policy, or existing U.S. Central Command (CENTCOM) operational orders and guidance, instead it is a complimentary supplement to them.

This document is based upon the best information available at the time of publication. It is not intended to define a standard of care and should not be construed as one. Neither should it be interpreted as prescribing an exclusive course of management. Variations in practice are inevitable and appropriately occur when clinicians take into account the needs of individual patients, available resources and limitations unique to an institution or type of practice.

VERSION 3.0, 23 APR 2021

CONTRIBUTORS

Leads: Maj Mary Stuever, MAJ Ashley Hydrick, Dr. Patricia Meza, COL Tamara Funari

CDR Stephanie Bedzis	Dr. Sean Keenan
Col. J. Scott Calder	COL Donald Kimbler
CDR R. Christopher Call	Ms. Linda Martinez
Maj Rebecca Castañeda	Ms. Lisa McFarlan
COL Amal Chatila	CPT Rachel Patrick
Lt Col Vicki Charbonneau	MSG Michael Remley
LTC Sally Delvecchio	LTC Johnnie Robbins
CDR Bambi Pish-Derr	Ms. Laura Runyan
Ms. Patricia Drouillard	Dr. Elizabeth Mann-Salinas
CDR Nichole Dutton	Mr. Phil Sartin
LTC Jennifer Edmundson	Col Stacy Shackelford
Dr. Erin Eickhoff	LTC Tom Sherbert
LTC Jennifer Gomes	Mr. Jim Sjovall
Lt Col Andrew Hall	Dr. Kimberly Smith
LTC Fred Hauser	Lt Col Steffanie Solberg
Mr. C. Ray Huntsinger	

TABLE OF CONTENTS

EXECUTIVE SUMMARY	1
ADDITIONS & REVISIONS	1
HOW TO USE	1
KEY DEFINITIONS	1
ACRONYMS AND ABBREVIATIONS	3
OPERATIONAL PLANNING	4
COMMAND AND CONTROL	4
COMMUNICATIONS	5
DOCUMENTATION	6
SUSPECTED, CONFIRMED CASES	6
PERFORMANCE IMPROVEMENT (PI)	6
TRANSPORT	7
PERSONAL PROTECTIVE EQUIPMENT (PPE) CONSIDERATIONS	8
FACILITIES MANAGEMENT	8
ISOLATION FACILITIES (ISOFACS)	9
QUARANTINE FACILITIES	10
SURGE CAPABILITIES	11
CRISIS STANDARDS OF CARE (CSOC)	11
CRITICAL RESOURCE CONSERVATION	12
ETHICAL CONSIDERATIONS	13
INFECTION PREVENTION AND CONTROL	14
TRIAGE/SECURITY	14
TRIAGE PROTOCOL AND EXAMPLE	14
INFECTION PREVENTION AND CONTROL MEDICAL CONSIDERATIONS	16
INFECTION PREVENTION AND CONTROL NON-MEDICAL CONSIDERATIONS	17
HUMAN RESOURCES	18
STAFF (PATIENT STAFFING RATIOS, RESILIENCE/WEILL-BEING)	18

SKILLS BUILDING/TRAINING.....	19
STAFF SURVEILLANCE AND STAFFING DECISIONS	21
TESTING	22
TYPES OF COVID TESTS.....	22
COVID TESTING STRATEGIES	22
SAMPLE POOLING.....	23
DoD TESTING ALGORITHMS	24
MEDICAL MANAGEMENT.....	24
IMMEDIATE CARE.....	24
PROLONGED CARE	24
DISCHARGE	24
TELEMEDICINE	25
BEHAVIORAL HEALTH	27
ARMED FORCES MEDICAL EXAMINER SYSTEM AND MORTUARY AFFAIRS	28
ANCILLARY SERVICES	29
PHARMACY	29
RADIOLOGY.....	29
LABORATORY (SCREENING/TESTING).....	29
ENVIRONMENTAL SERVICES/HOUSEKEEPING.....	29
FORCE HEALTH PROTECTION.....	30
FIELD SANITATION AND ISOLATION.....	30
FOOD SAFETY AND FOOD SERVICE	31
VETERINARY CARE	31
ANIMAL TO HUMAN TRANSMISSION	32
CONSIDERATIONS FOR ANIMAL CARE PERSONNEL	32
SUPPLIES	33
POINTS OF CONTACT	34
INDEX.....	34
APPENDIX A: PANDEMIC RESPONSE PLANNING CHECKLIST	36
APPENDIX B: ADDITIONAL TRAINING AND RESOURCES	38
APPENDIX C: MITIGATING TEAM STRESS	39
APPENDIX D: SLEEP CHECKLIST.....	40
APPENDIX E: COVID-19 IN FORWARD-DEPLOYED SETTINGS EXECUTIVE SUMMARY.....	42

LEGEND OF FIGURES & TABLES

Figure 1. Good, Better, Best Characteristics of PPE.....	8
Table 1. Ethical Values to Guide Rationing of Absolutely Scarce Healthcare Resources in a COVID-19 Pandemic....	12
Figure 2. Paradigms for Changing Standards of Care (NAM 2009).....	13
Figure 3. Patient Receiving and Triage Protocol.....	15
Table 2. Quarantine versus Isolation	16
Figure 4. Tiered Staffing Strategy	18
Table 3. COVID-ICU	19
Table 4. COVID-ICW	19
Table 5. CDC Covid Testing Strategies	23
Table 6. Local USCENTCOM Advisors For Urgent Needs	26
Table 7. MEDCEN Contact List	26
Figure 5. Clinician's Guide of Mobile Health Apps.....	28
Figure 6. PPE Recommendations for the MHS.....	33

EXECUTIVE SUMMARY

This playbook is aimed at healthcare, public health providers, and staff in the United States Central Command (USCENTCOM) Area of Responsibility (AOR). The USCENTCOM AOR is a medically austere operational environment where Role 3 (R3) is the highest level of medical capability. All planning assumptions proceed from there. This playbook is a concise source of key information, references, and materials for providers and healthcare leaders to easily review. The playbook is comprehensive but **not all inclusive**, and does not negate the need for communication and information sharing with local, Joint Task Force (JTF), Service Component and USCENTCOM Headquarters (HQ) Command and Surgeon Cells.

ADDITIONS & REVISIONS

Revisions and updates were completed throughout this document to be current as of 23 Apr 2021. Major additions and revisions include:

- Updated information on guidelines for Force Health Protection, to include: isolation and quarantine, including discharge and return to duty guidelines
- Added information on testing for general screening and diagnostic testing
- Added information on the new COVID vaccination
- Added information on the treatments for COVID
- Added updates for patient tracking in theatre
- All other revisions involve addition of resource links throughout and minor refinement of language for clarity and grammar

HOW TO USE

This document provides key information in text and graphic images for rapid review. The Table of Contents will link users to the topic they want to review, which will contain key facts and figures, as well as linking users to external resources that can provide greater detail. Users are encouraged to use this playbook as a tool to connect to best practice resources that will augment existing clinical and military expertise and local, JTF, Service Component, and USCENTCOM guidance.

KEY DEFINITIONS

1. **AE Patient Classification:** A basic metric to convey patient acuity and level of infectivity to others. Patient classification will be performed in accordance with the *Department of the Air Force Instruction DAFI 48-107, EN Route Care Documentation, Volume 3, 17 Dec 2020*.
2. **Airborne Spread:** Spread of disease via small liquid particles (aerosols) that remain aloft for prolonged periods of time and may travel longer distances. Airborne precautions aim to mitigate this method of transmission.
3. **Antibody:** Proteins produced by infected individual as part of their immune response to infection by the pathogen.

4. **Antigen:** Proteins or other cellular or chemical features associated with the pathogen (e.g., viral DNA, surface proteins).
5. **Close Contact:** Close contact is defined by the Centers for Disease Control and Prevention (CDC) as someone who was within 6 feet of an infected person for a cumulative total of 15 minutes or more over a 24-hour period starting from 2 days before illness onset (or, for asymptomatic patients, 2 days prior to test specimen collection) until the time the patient is isolated.
6. **Contact Spread:** Spread of disease via direct contact with an infected patient or contaminated surface. Contact Precautions aim to mitigate this method of transmission.
7. **Droplet Spread:** Spread of disease via relatively large liquid particles that settle from the air quickly (within a few feet). Droplet Precautions aim to mitigate this method of transmission.
8. **Intermediate Care Ward (ICW):** An inpatient nursing unit that typically accommodates patients requiring med surgical to progressive (step-down) – level care. In pandemic conditions, this definition may broaden to incorporate levels of care similar to mass casualty. Staffing considerations and Knowledge, Skills and Abilities (KSAs) will be commensurate with the roles required. The mission may require staff to increase their KSAs. Additional considerations may be needed for patients requiring isolation. In a crisis environment, all staff are asked to work at the top of their license.
9. **Isolation:** The separation of an individual or group infected or reasonably believed to be infected with a communicable disease from those who are healthy in such a place and manner to prevent the spread of the communicable disease. Isolation is a medical term and ordered by a medical provider, but requires command support for successful execution.
10. **Pathogen:** an organism that causes disease (e.g. virus, bacteria, fungus, or parasite). The term is also used to describe prions – non-living protein particles that display infectious behavior.
11. **Patient Under Investigation (PUI):** A patient with signs and symptoms consistent with known possible presentations of COVID-19 with potential exposure to the virus. Potential exposure to the virus is defined as close contact with known or other suspected cases and/or travel through regions with widespread sustained transmission of COVID-19. In areas where COVID-19 is already widespread, symptoms alone may make the diagnosis of “PUI.” A PUI may refer to: 1) an individual whose diagnostic testing has not yet been performed, 2) an individual with symptoms consistent with COVID-19 but diagnostic testing is unavailable, or 3) an individual with high index of suspicion for COVID-19 whose initial COVID test was negative. All PUIs must be isolated. All patients being admitted to the hospital for suspected COVID are considered PUIs until tested for COVID.
12. **Quarantine:** The separation of an individual or group that has been potentially exposed to a communicable disease, but is not yet ill, from others who have not been so exposed, in such a manner and place to prevent the possible spread of the communicable disease. This is a form of Restriction of Movement (ROM) and is a command function that is medically supported. Quarantine is a commander’s responsibility.
13. **Social Distancing:** The practice of reducing close contact between people to slow the spread of disease. It includes: wearing masks that cover the nose and mouth, limiting large group gatherings (no more than 10 persons), closing buildings and gathering spaces, cancelling events, and advising people to stay six feet apart as much as possible. Preparation of isolation berthing, including activities of daily living (e.g., hydration, food, hygiene, trash disposal) is addressed on [pages 25-26 of the COVID-19 CPG for Austere Operational Environments](#).

ACRONYMS AND ABBREVIATIONS

AE - Aeromedical Evacuation
AFMES - Air Force Medical Examiner System
AOR - Area of responsibility
C2 - Command and Control
CENTREX COVID tracker: using this system to track patients.
CJTF - Combined Joint Task Force
CCIR - Commander's Critical Information Reporting
CCOP - Central Command Operating Procedures
CCSG - Command Surgeon
CHU - Containerized Housing Unit
CONOPS - Concept of Operations
CONPLAN - Concept of Operations Plan
CSOC - Crisis Standards of Care
DHA - Defense Health Agency
EXORD - Executive Order
FFR - Filtering Facepiece Respirators
HIPAA - Health Insurance Portability and Accountability Act
HPCON - Health Protection Conditions
ICW - Intermediate Care Ward
KSA - Knowledge, Skills and Abilities
MS - Medication Safety
MSAT - Medical Situational Awareness in the Theater
MWR - Morale Welfare and Recreation
PHEO - Public Health Emergency Officer
PHEM - Public Health Emergency Management
PFA - Psychological First Aid
PMG - Practice Management Guide
PPE - Personal Protective Equipment
PS - Patient Safety
PUI - Person Under Investigation
QS - Quality and Safety
SIPRNet - Secret Internet Protocol Router Network
SME - Subject matter expert
TPMRC-E - Theater Patient Movement Requirements Center East
USCENTCOM - United States Central Command
USTRANSCOM - U.S. Transportation Command
VITAL-T - Virtual Inspection and LINKUP in Theater
VH - Virtual Health

OPERATIONAL PLANNING

The USCENTCOM CONPLAN 1251-15, Regional Concept Plan for Preparation and Response for Pandemic Influenza and Infectious Disease (PI&ID), delineates the policies, actions and requirements for the employment of military resources within the USCENTCOM AOR for PI&ID preparation and response plan, along with other plans, are available for download on the [USCENTCOM SIPRnet website](#). You can also reach the SIPRNet through <https://ccsg.nonrel.centcom.smil.mil/Private/SitePages/Home.aspx> and click on the COVID-19 link. There will be links to EXORDS and instructions. Refer to CCJ3 for assistance if copies of instructions are needed.

Specific Health Service Guidance is delineated in the Annex Q to the *USCENTCOM EXORD Novel Coronavirus Outbreak Response Operations* in the UCENTCOM SIPRNet link.

An excellent summary of medical operational experiences was authored by Task Force Medical (TFM) 14. The Memorandum for record: Role 3 Standard Operating Procedure (SOP) for Coronavirus (COVID-19) Triage and Admission Criteria, 18 Aug 2020, provides guidelines for TFM 14 Role 3 responsibilities while caring for suspected COVID-19 positive patients.

COMMAND AND CONTROL

Coordination is essential between Command Staff, medical teams, and public health/preventive medicine assets. Given the highly complex nature of the COVID-19 disease and need for subject matter expertise (SME), Commanders are encouraged to designate a COVID-19 response and planning team consisting of medical and public health/preventive medicine SMEs along with operational planners. Task Force Med Leaders are encouraged to coordinate with respective Surgeon teams and Force Health Protection assets to translate and communicate all guidance and policy down to the most forward units with attention to operational security and patient privacy laws. In order to facilitate a coherent Joint effort, the response should follow established DoD procedures as outlined in [DODI 6200.03 Public Health Emergency Management \(PHEM\) Within the DoD, 28 Mar 2019](#) and [DODI 6055.17 DoD Emergency Management \(EM\) Program, Change 3 effective 12 Jun 2019](#).

All leaders should operate under a number of considerations/assumptions including (but not limited to):

1. Command and Control (C2) authorities will remain unchanged unless otherwise directed. Combined Joint Task Force Surgeon (CJTF) and Component Surgeon cells should continue to collaborate and communicate closely with higher headquarter authority and disseminate information to the most forward units.
2. Current USCENTCOM missions will continue, unless otherwise directed, throughout the COVID-19 pandemic, daily tasks may need to be reprioritized as required by a pandemic response to maintain surgical and critical care capabilities while minimizing spread of infectious disease in theater.
3. Leaders are encouraged to collaborate and delegate authority to improve comprehensiveness planning and operation activities. Subordinate leaders should be empowered to utilize their expertise and innovation within the boundaries set-forth by pre-established policy and guidance. The USCENTCOM Health Protection Condition (HPCON) Checklist, Joint Force Health

Protection Team, COVID-19 Crisis Action Team (when available) will assist appropriate actions listed in [Appendix A](#).

4. Communication is key for interdisciplinary COVID-19 planning and response. Limit jargon and clearly define all acronyms and unfamiliar terms, for example. Commanders should communicate key information to subordinate units.
5. Logistical and patient movement channels are likely to be altered by the pandemic both in theater and at receiving Role 4 Contiguous United States (CONUS) and Outside the Contiguous United States (OCONUS) facilities. These challenges must be overcome by clear communication and adaptive planning strategies.

RESOURCES

Refer to Headquarters, CJTF, and Component guidance for specific information and requirements with regards to operational planning guidance.

The U.S. Department of Health and Human Services provides [Critical Care Planning-COVID-19 Quick Notes](#) a two-page document which describes operationalization of the concept in three major categories: space, staff, supplies, and provision of critical care.

Planning and Preparation, pg6, Implications of COVID-19 on Surgical Care, p8, [DoD COVID-19 PMG v7.0, 03 Mar 2021](#).

[COVID-19 Response and Prevention Planning-Knovel \(Elsevier\)](#)

[DODI 6200.03 Public Health Emergency Management \(PHEM\) Within the DoD, 28 Mar 2019](#)

[DODI 6055.17 DoD Emergency Management \(EM\) Program, Change 3 effective 12 Jun 2019](#)

COMMUNICATIONS

All USCENTCOM bases and facilities should establish a local and regional PACE (Primary-Alternate-Contingency- Emergency) plan for both operational and clinical communication incorporating social distancing and division of labor during the pandemic response period.

The USCENTCOM Component and CJTF teams may have prospectively published local and regional PACE plans for both operational and clinical consultation and communication. Forward-stationed medical teams/medics should identify and test these options PRIOR to needing urgent consultation.

Division of labor and social distancing could strain all routine secure and unsecure DoD communication and collaboration platforms (i.e. teleconferencing video conferencing networks, remote access email), resulting in the incorporation of non-DoD unsecure platforms (e.g., Zoom, Skype, WhatsApp).

Operational security and patient privacy must be a primary consideration when selecting communication platforms, especially when using non-DoD platforms.

UPDATE 2.0: *In answer to the social distancing challenge in the work place, the DoD has established an approved virtual communication platform to improve personnel connection and information sharing. This platform is approved for controlled unclassified information sharing and is noted HIPAA compliant. It connects with individuals' regular Non-classified Internet Protocol Router (NIPR) email accounts for ease of communications and scheduling. For more information on how to get connected visit the following link through June 2021: <https://www.cloud.mil/CVR/>.*

RESOURCE

Refer to the [Planning and Preparation, p12, DoD COVID-19 PMG v7.0, 3 March 2021](#)

DOCUMENTATION

Documentation of patient care and movement should continue via the usual platforms (i.e. paper or electronic charts) as previously established at the local treatment facility. The appropriate ICD-10 codes, and symptoms (e.g. fever, cough and shortness of breath), should be entered as detailed as possible in order to capture these patients in Theater Medical Data Store (TMDS), Medical Situational Awareness in the Theater (MSAT) and Medical Combined Operating Picture (MEDCOP) and future databases and overall future performance improvement opportunities.

JTS has published the ICD-9 and ICD-10 Codes to accompany the new COVID-19 registry under development. The codes can be found on the following JTS website:

https://jts.amedd.army.mil/assets/docs/education/COVID-19_ICD-9_ICD-10_Codes.pdf

Please ensure that outpatient encounters are closed at the end of the patient encounter (i.e. when the patient departs medical facility).

SUSPECTED, CONFIRMED CASES

All patients, staff, and support personnel with symptoms (fever, cough, shortness of breath) should be tested for COVID-19 using the available confirmatory diagnostic test.

All persons who test positive need to be moved into isolation and if medical management is required, considered for evacuation to OCONUS or CONUS locations as designated for the evacuation plan per that region (follow the guidance provided in the Transport section of this Playbook).

Command and medical teams are responsible for establishing a plan for patient tracking and re-unification (for family notification of patient status) locally. In accordance with USCENTCOM regulations, local Commanders will include COVID-19 patient tracking as part of their Commander's Critical Information Reporting (CCIR). Patient tracking for the USCENTCOM AOR will be accomplished using SIPR COVID-19 CAT SharePoint/Trackers & Updates/Patient Trackers by Component.

PERFORMANCE IMPROVEMENT (PI)

The JTS has partnered with the Uniformed Services University of Health Sciences (USU) to develop a registry for COVID-19 patient data acquisition. Detailed documentation including appropriate ICD-10 codes are essential to assist key term triggers for audits. If using hard copy/paper charting, all documentation should be uploaded to Theater Medical Data Store as soon as possible.

The JTS COVID registry will ease tracking and monitoring the progress of this disease process, while evaluating the quality and possibility for improvement of delivery of care. In the meantime, units should designate a person responsible for tracking and monitoring the COVID-19 patient care and patient movement. The JTS COVID-19 Registry is the only approved data repository for CENTCOM.

For information email: DHA.JBSA.j-3.List.JTS-PIP@mail.mil (JTS PI team).

The following list provides the intermediate and after action reviews that provide important sources for sharing best practices and innovative ideas to enhance performance improvement:

1. [COVID-19 Monitoring and Response Among U.S. Air Force Basic Military Trainees — Texas, March–April 2020, MMWR, 05 June 2020.](#)
2. [COVID-19 in Erbil Air Base: Lessons Learned Managing Multiple COVID-19 Outbreaks at a Deployed, Austere Location](#) (See Appendix E: [COVID-19 in Erbil Air Base Executive Summary](#)).

TRANSPORT

Transport guidance has changed frequently and is ever changing with the current environment. Check with local Operations cell to ensure most current guidance is followed. As of the date of this publication:

- All patients with symptoms severe enough to require admission to a medical facility for symptom management and/or supplemental oxygen should be considered for evacuation.
- Host Nation admitted patients may remain in hospital at commander's discretion.
- Navy Afloat remains on ship unless ship's capability is in danger of being exceeded (either by volume or the severity of the patient.)
- Asymptomatic COVID+ or mildly symptomatic patients should stay but a commander may decide to evacuate.

For patients with a new (+) COVID test within 72 hours of patient transport by AF Aeromedical Evacuation (AE/CCATT) or civilian air ambulance, in addition to usual contact tracing procedures, please notify the regional TRANSCOM Patient Movement Requirements Center (TPMRC) that services that MTF (TPMRC-A DSN (312) 779-4200, TPMRC-E DSN (314) 480-8040, TPMRC-W DSN (315) 448-1609); the U.S. Transportation Command (USTRANSCOM) Instruction 41-02, 11 July 2019 outlines Patient Movement of Contagious and potentially exposed casualties. This playbook is a summary that pertains specifically to COVID-19. ***The guidance for movement and evacuation of COVID-19 positive patients is rapidly evolving. Early and close coordination of these movement requests with Theater Patient Movement Requirements Center East (TPMRC-E) and next higher headquarters is crucial.***

To ensure correct resource allocation and transport prioritization, units should refrain from using locally derived patient categorization. Patient classification should remain IAW *DAFI 48-107, EN ROUTE CARE DOCUMENTATION, Volume 3, 17 December 2020.*

Movement of multiple COVID-19 patients will be performed using a DoD certified and approved contagious patient movement system. Planning considerations must take into account downtime for decontamination of aircraft following COVID-19 patient evacuation. Upon consultation and appropriate approvals, capabilities developed outside of the DoD for moving patients exposed to or infected with COVID-19 may be used for approved missions via commercial aircraft. The sending MTF must coordinate movement with TPMRC-E for both intra and inter-theater PM.

Refer to local, CJTF, Component, and USCENTCOM guidance for specific knowledge on Operational Planning assumptions, considerations and guidance located on the [USCENTCOM Surgeon \(CCSG\) SharePoint](#).

COVID-19 is a respiratory virus that requires up to advanced droplet (droplet/airborne) precautions depending on the risk of activity (below). All re-use and extension of PPE MUST be done in accordance with CDC, JTS, and USCENTCOM best practices and guidance.

Good	Better	Best	Best of the Best
Standard + Droplet Precautions *Don and doff procedures occur in warm areas –Gloves-single pair –Scarf or Balaclava –Eye Pro or Goggles –Disposable T-shirt	Droplet + Airborne Precautions *Don and doff procedures occur in warm areas –Double Gloves –Surgical Mask –Eye Pro or Goggles –Disposable Gown	Droplet + Airborne Precautions *Don and doff procedures occur in warm areas –Double Gloves –N95 Mask –Full Face Shield –Disposable Gown –Hood or Head Cover	Droplet + Airborne Precautions **High risk exposure to aerosolized droplets** –Double Gloves –Surgical Mask covering –N95 Mask –Full Face Shield –Tyvek* Suit - Hooded –Disposable Foot Covering

◆ ————— ◆ Intubation

Relative Risk / Intensity of Exposure

LOW —————> HIGH

RESOURCES:

[USCENTCOM Infection Prevention and Control Policy in a Deployed Setting](#) or service component site such <https://portal.arcent.swa.army.mil/covid19opt/SitePages/Home.aspx>

- [CDC Decontamination and Reuse of Filtering Facepiece Respirators](#)
- [CDC Strategies to Optimize the Supply of PPE and Equipment](#)

Employed cleaning protocols should ensure adequate sanitization in all environments, including quarantine/isolation/patient care areas, as well as all workspaces and quarters. All non-dedicated, non-disposable patient care medical equipment should be cleaned and disinfected according to manufacturer's instructions and facility policies. The U.S. Environmental Protection Agency (EPA) has compiled a list of [Disinfectants for Use against SARS-CoV-2 \(COVID-19\)](#).

8

Opening high-flow doorways in hallways can reduce the number of high-touch surfaces in a facility, while closing the doors of individual office spaces can reduce cross-contamination of virus across office spaces. Doorway management approaches MUST consider operational security, facility security, safety, and privacy guidelines.

- **Entrances/Exits:** Hand sanitizer dispensers should be placed near entryways. Limiting the flow of traffic and personnel in and out of high traffic areas should be considered to reduce cross-contamination.
- **Bathroom Entrances/Exits:** Hand washing signs should be clearly posted. When possible, position trash cans inside bathrooms near the door to allow paper towel use and disposal for no-touch exit.
- **Single occupancy:** Best practice is to isolate personnel and patients in single occupancy to reduce the spread. Areas that were once open bays should be evaluated for ability to convert to single occupancy, if possible.
- **Negative pressure rooms:** The following are the best practices to limit the exposure of patients and staff: check with facilities management to ensure that the negative pressure rooms are functioning, evaluate for the possibility of adding negative pressure rooms as needed for the facility.

Note: *Separate ventilation is important.*

RESOURCES:

[USCENTCOM Infection Prevention and Control Policy in a Deployed Setting \(CCOP-02\)](#)

[CDC Cleaning and Disinfecting your Facility](#)

[CDC Cleaning and Disinfection for Households](#)

ISOLATION FACILITIES (ISOFACs)

Isolation facilities (ISOFACs) will need to be established in accordance with best practice and other applicable guidance while taking into consideration the following issues:

- While both COVID-19 positive and PUIs require isolation, it is important not to mix them to prevent the spread of COVID-19 to a non-infected PUI.
- 24/7 communication between COVID-19 Isolation berthing and the medical personnel is mandatory as patients may develop worsening symptoms.
- Design ISOFAC policies and areas to allow maximum autonomy and self-management by the ISOFAC occupants and their units, including cleaning, food delivery, and transport.
- A designated “dead drop” area allows unit representatives to deliver food and supplies while avoiding inadvertent interactions with COVID positive patients.
- Identify a camp mayor for each isolation facility. Camp mayors should become familiar with base policies and instructions for requested services, repairs, and maintenance.
- Isolation facilities must include plans for medical or security evacuations including a designated shelter area.

ISOFACs are designed to allow outpatient recovery similar to a COVID positive patient recovering at home in CONUS. Unfortunately, most forward-deployed locations berth individuals in group berthing, this arrangement makes isolation difficult to achieve. While individual berthing with separated bathrooms is optimal, patient isolation cohorts may be the only option available. Isolation berthing should be designed to allow adequate comfort for multiple patients of varying symptom severity. Isolation areas should be well demarcated (e.g., “Isolation Area - No Unauthorized Access!”) and include posted guides on proper PPE procedures. Additionally, consider any medical evacuation routes in the event of worsening COVID-19 disease.

If possible, ISOFACs should include an entry anteroom that separates the outside from the isolation rooms and may include a designated “clean area” and “dirty area.” The clean area should be stocked with PPE and visual donning instructions. The “dirty area” should include a biohazard disposal bin and hand sanitizing station for doffing PPE.

A designated bunker should be pre-identified for the ISOFAC occupants in the event of an indirect fire. To mitigate transmission to non-isolated individuals, consider taping PPE bags to each door for patients to grab on their way to the bunker or other quick access methods.

Some ISOFACs may allow for individual/paired isolation (e.g. Containerized Housing Unit [CHU]), whereas others may require cohort isolation (e.g. tent). Subsequently, individual/paired isolation areas should be considered the preferred isolation site for females and individual cases. For unit outbreaks, cohort isolation may result in improved unit morale and accountability. Cohort isolation of confirmed cases can join pre-existing cohorts, whereas presumed positives (i.e. PUIs) should be isolated individually.

Medical personnel should conduct a daily sick call and should wear full PPE during encounters. Common symptoms may be immediately treated with over-the-counter medications; any individual with specific concerns should be evaluated. For worsening symptoms, consider obtaining vital signs and a NEWS score. Refer to [Screening and Triage: Early Recognition of Patients with COVID-19 in the DoD Practice Management Guideline](#) for current best practices.

QUARANTINE FACILITIES

Many sites have established a centralized base quarantine facility. However, it is important to realize that during a significant outbreak, the number of close contacts can quickly overwhelm any centralized facility. Therefore, units should have pre-established plans to quarantine close contacts within their living spaces. Quarantine is a command function, although medical issues that arise during quarantine should be referred to the appropriate medical unit.

Consider the following issues when establishing a comprehensive quarantine plan:

- Quarantine is a unit responsibility, though medical experts have much to offer in advisement and assistance for quarantine planning, tracking, and execution.
- Establishing a base quarantine camp requires strong partnerships between medical experts, Base Operating Support Integrator (BOS-I), Base Defense Operation Center (BDOC), and base leadership.

- Designating one or two days per week to initiate travel quarantine cohorts helps reduce limited quarantine resources and streamlines throughput.
- Base quarantine camps require daily walkthroughs from a Camp Mayor to mitigate current or future concerns. Daily vital sign checks for quarantined personnel increases the risk of close contact exposure with little benefit.

At a minimum, quarantine plans must include designated berthing areas, latrines, and showers for close contacts that are separated from the general populace. Meals should be delivered to quarantined individuals. Examples of unit quarantine plans include: 1) quarantining close contacts in their CHU rooms with a designated quarantine latrine, 2) moving close contacts into a designated quarantine tent that includes adjoining chemical latrines and portable shower units, and 3) deploying temporary fencing around an affected tent to designate it as a quarantined tent. Medical teams may provide guidance as units establish their quarantine plans and facilities.

Bases that serve as theater gateways experience a high volume of travel quarantine requirements. Quarantine cohorts must start and end their quarantine period as a group. However, it may prove infeasible to establish travel quarantine cohorts for each day's arrivals. A base may choose to designate a fixed day(s) of the week for travel quarantines to begin. As an example, a base may begin a 14-day travel quarantine cohort every Tuesday. Incoming personnel who arrive Wednesday through Monday are placed into the upcoming quarantine cohort upon arrival; however, the 14-day quarantine period does not start until Tuesday. The entire cohort is then released to their respective units two weeks later. Individuals who do not abide by the quarantine order restart their 14-day quarantine along with all others in their cohort. This model requires three quarantine areas that rotate between the "fill" cohort, a cohort in their first quarantine week, and a cohort in their second quarantine week. Privacy dividers (e.g., central tent dividers) can allow males and females from the same cohort to share a cohort area.

SURGE CAPABILITIES

Units will investigate and proactively plan inpatient and outpatient surge capabilities within their existing resources. Special consideration should be given to preventing cross-infection of the patient population. A sample checklist to assist with surge capability planning at the role 1, 2, 3 level is at [Appendix A](#).

CRISIS STANDARDS OF CARE (CSOC)

CSOC are guidelines that are applied when a pervasive or catastrophic disaster makes it impossible to achieve the usual standards of care.

The National Academies Press released an article from the National Academies of Sciences, Engineering and Medicine entitled, [Rapid Expert Consultation on Crisis Standards of Care for the COVID-19 Pandemic](#). This expert guide assists the provider in establishing a rationale for the implementation of crisis standards of care.

Additional assistance in developing and implementing this crisis standards of care can be accessed in the milBook through the following link: <https://www.milsuite.mil/book/groups/covid-19-clinical-operations-group>.

CRITICAL RESOURCE CONSERVATION

There are limited resources for both the treatment and the diagnosis of COVID-19. Within the framework of responding to a global crisis, there must be evaluation and conservation of critical resources in terms of personnel, supplies and equipment. This section offers the resources for establishing a protocol for resource management with the focus on conservation. This is a broad approach and should be tailored to the resources of the facility for which the provider is managing.

RESOURCES:

[USCENTCOM Policy for the Decontamination and Reuse of Filtering Facepiece Respirators \(FFR\) Such as the N95](#) (CAC required)


[CDC Decontamination and Reuse of Filtering Facepiece Respirators](#)

[CDC Strategies to Optimize the Supply of PPE and Equipment](#)

Table 1 published by the *New England Journal of Medicine* is a community standard consensus opinion of values.

Table 1. Ethical Values to Guide Rationing of Absolutely Scarce Healthcare Resources in a COVID-19 Pandemic

Ethical values & guiding principles	Application to COVID-19 pandemic
Maximize benefits	
<ul style="list-style-type: none"> Save the most lives Save the most life-years – maximize prognosis 	<ul style="list-style-type: none"> Receives the highest priority Receives the highest priority
Treat people equally	
<ul style="list-style-type: none"> First come, first serve Random selection 	<ul style="list-style-type: none"> Should not be used Used for selecting among patients with similar prognosis
Promote and reward instrumental value (benefit to others)	
<ul style="list-style-type: none"> Retrospective – priority to those who have made relevant contributions Prospective – priority to those who are likely to make relevant contributions 	<ul style="list-style-type: none"> Gives priority to research participants and healthcare workers when other factors such as maximizing benefits are equal Gives priority to healthcare workers
Give priority to the worst off	
<ul style="list-style-type: none"> Sickest first Youngest first 	<ul style="list-style-type: none"> Used when it aligns with maximizing benefits Used when it aligns with maximizing benefits such as preventing spread of the virus.

Figure 2. Paradigms for Changing Standards of Care (NAM 2009)

- Unless temporary, requires state empowerment, clinical guidance, and protection for triage decisions and authorization for alternate care sites/techniques. Once situational awareness is achieved, triage decisions should be as systematic and integrated into institutional process, review, and documentation as possible.
- Institutions consider impact on the community of resource use (consider “greatest good” versus individual patient needs—e.g., conserve resources when possible), but patient-centered decision-making is still the focus.
- Institutions (and providers) must make triage decisions—balancing the availability of resources to others and the individual patient’s needs—shift to community-centered decision-making.

ETHICAL CONSIDERATIONS

During a local, regional, national and global crisis, many decisions may require ethical considerations. These decisions are best managed in a multidisciplinary/multi-level command structure setting employing a good, better, best approach. See the following Resources listed.

RESOURCES:

[DoD COVID-19 PMG v7.0, 3 March 2021:](#)

- Ethics of Clinical Research during a Pandemic, p50
- Palliative Medicine during the COVID-19 Pandemic, p81-85
- Adjunctive Therapies: Treatment Protocols p49
- Planning And Preparation, p90-93
- [Crisis Standards of Care: A Toolkit for indicators and Triggers](#), p138; Board on Health Sciences Policy; Institute of Medicine

INFECTION PREVENTION AND CONTROL

TRIAGE/SECURITY

Medical triage is a familiar concept for all military medical personnel. The primary goal of saving as many lives as possible remains the same for pandemic disease triage. However, the operational concepts shift to the following main ideas:

1. Identifying and cohorting infectious or potentially infectious persons as soon as possible.
2. Identifying infected individuals who will most likely deteriorate rapidly/require prolonged advanced medical care to prioritize medical evacuation.
3. **During the triage process, treat patients as if they are possibly infected, and screen for appropriate need to continue to isolate patients based on symptoms.**

RESOURCES:

[Screening and Triage: Early Recognition of Patients with COVID-19, p11](#)

[Appendix B: Example Triage Protocols during COVID-19 Pandemic, p96](#)

[Management of COVID-19 in Austere Operational Environments CPG v3.0, 23 Apr 2021](#)

[Actions on Identification of a COVID-19 PUI, p5](#)

[Providing Medical Care to a COVID-19, p13](#)

[CDC Information on Contact Tracing and Epidemiologic Interviewing](#)

[Pandemic Influenza Preparedness and Response Guidance for Healthcare Workers and Healthcare Employers](#)

Complete DD Form 3112, Personnel Accountability and Assessment Notification for Coronavirus Disease 2019 (COVID-19) Exposure, April 2020. Turn in [DD Form 3112](#) to Public Health.


TRIAGE PROTOCOL AND EXAMPLE

The following is an example of the patient intake and triage approach used by Task Force Medical 14:

- The trauma and COVID-19 screening and treatment locations were separated to minimize exposure.
- A speaker phone was placed outside the hospital entrance for direct access to the Patient Administration Department (PAD) line.
- Patients answered a series of questions to dictate their course.
- If COVID-19 was suspected and the patient was stable, they would wait to be assessed in the COVID-19 screening trailer located 50 meters from the hospital entrance.
- For patients deemed unstable, the evaluation would be conducted within the COVID-19 ICU.
- Pending the results of the provider's assessment, the patient would either go into quarantine in the respective base location or get admitted to the COVID-19 portion of the medical footprint.
- Those that were able to be quarantined were evaluated at various intervals and provided meals by the hospital staff until release criteria was met.

Refer to Figure 3. Patient Receiving and Triage Protocol for more details about triage protocol.

Figure 3. Patient Receiving and Triage Protocol



*Image Adjusted from the USAG West Point COVID-19 Response, West Point Community Playbook and JTS Overview of COVID-19 in Austere Environments Infographic

INFECTION PREVENTION AND CONTROL MEDICAL CONSIDERATIONS

Standard hospital infection prevention and control procedures will be implemented immediately in accordance with practice guidelines and best practice principles. Isolation (aka Transmission-Based Precautions) and Quarantine protocols are “step above” infection prevention and control measures employed in an epidemic/pandemic situation.

RESOURCES:

[Routine Infection Control and Prevention Principles: USCENTCOM Infection Prevention and Control Policy in a Deployed Setting](#)

[CDC Infection Control Guidelines & Guidance Library](#)

[Management of COVID-19 in Austere Operational Environments, Quarantine and Isolation Procedures, 23 Apr 2021, p8](#)

[Framework for Healthcare Systems Providing Non-COVID-19 Clinical Care during the COVID-19 Pandemic](#)

Table 2. Quarantine versus Isolation

	Quarantine	Isolation
Whose activity?	Command approves and tracks quarantine activities. A person does not have to be seen by a health provider unless symptoms occur in quarantine.	A person must be seen by a healthcare official or provider to enter isolation.
What is it?	The separation of an individual or group that has been potentially exposed to a communicable disease, but not yet ill to prevent spread of disease.	The separation of people who are infected or with suspected infections with a communicable disease to prevent spread of that illness.
Who is it for?	Potentially exposed persons to COVID-19 + individual OR PUI (<i>Per definition, a PUI has symptoms and requires isolation, not quarantine</i>).	Known COVID-19+ or suspected COVID-19 cases as determined by a healthcare official or provider.
How long is it?	14 days after the last exposure, assuming all release criteria is met. Release should be coordinated with a designated COVID-19 provider. Follow current DoD/CENTCOM policy.	10 days and 48 hrs after resolution of symptoms or until released by designated healthcare provider
What does this mean for daily life?	Stay in quarters or designated quarantine areas. Limit/prohibit movement to essential common areas (e.g., dining facilities, laundry facilities, and bathroom/hygiene facilities) and duty location. Practice standard hygiene and precautionary measures. Wear a mask or face covering as directed if required to leave quarantine areas. Quarantined individual and teammates should monitor symptoms and report any changes to the supervisor and a healthcare provider.	Person CANNOT leave designated isolation location except in emergency or if instructed by medical personnel. Food, medication and other supplies MUST be delivered by designated personnel to the isolation area. Designated personnel will treat and monitor as determined by a health provider. Medical evacuation according to Command guidance.

INFECTION PREVENTION AND CONTROL NON-MEDICAL CONSIDERATIONS

Outside of medical settings, personnel **must** practice basic public health and infection prevention guidelines that focus on:

- Social Distancing
- Shelter-In-Place Activities
- Hygiene and General Protection Measures

Social Distancing is the practice of maintaining a minimum amount of separation between individuals and all other persons outside of their routine close contacts. Examples include roommates in barracks/quarters, family members in the same household, and teams that routinely experience unavoidable close contact in execution of daily duties (e.g., medical staff). For social distancing consider the following:

1. Ensure that all personnel can maintain at least 6-ft spacing between themselves at all times.
2. Ensure the lowest possible number of personnel in duty spaces (as needed to execute mission) and prevent (as much as possible) sharing of workspaces and equipment:
 - a. Reduce the number of personnel in duty areas by instituting teleworking, day-on/off or week-on/off.
 - b. Cohort all personnel into specific teams and ONLY persons from that team will work at the same time.
 - c. Hand-overs will be done remotely or with a limited person(s) from the departing team.
 - d. Each team cleans and disinfects the space, ESPECIALLY high touch surfaces prior to departing (See [Hygiene and General Protection Measures](#)).

Shelter-in-Place Restrictions require that all personnel limit their activity to quarters and their designated duty location (ONLY during their scheduled work period). This mandates NO social gathering or non-essential activities outside of personal areas AND closure of non-essential facilities (e.g., MWR, shoppettes). Essential activities/facilities include dining facilities (See [Food Safety and Food Service](#) section for more information duty activities as scheduled), medical care, and personal laundry.

Hygiene and General Protection Measures. All personnel should maintain an elevated awareness of personal hygiene and routine cleaning practices that will prevent the spread of the virus including hand hygiene, routine wear of face coverings outside of personal areas, daily personal hygiene and grooming, cleaning/disinfection of high touch surfaces and common areas.

RESOURCES:

[Non-Medical Infection Control, USCENTCOM Infection Prevention and Control Policy](#), p17-18. CAC required
[CDC How to Protect Yourself and Others](#)
[CDC Cleaning and Disinfecting Your Facility](#)
[CDC Cleaning and Disinfecting Your Home](#)
[OSHA Guidance on Preparing Workplaces for COVID-19](#)


HUMAN RESOURCES

STAFF (PATIENT STAFFING RATIOS, RESILIENCE/WEEL-BEING)

Outbreak response requires advanced staffing approaches to ensure safety and well-being of staff and patients, while limiting the spread of infectious disease. Facility managers should identify and assign essential personnel as per operational planning. Plans should identify tiered strategy planning and minimal risk approach. The tiered staffing strategy for pandemic (below) will be applied and tailored according to local resources and manning. Exact staffing plans and staffing ratios will depend on personnel, resources, and mission requirements. Additional considerations include:

1. Designated and/or specialized COVID-19 care team separate from combat operations.
2. Designated runners to ancillary services to reduce donning/doffing activities and protecting PPE supplies.
3. Assignment and training of non-medical personnel to assist as runners, computer and administrative activities (e.g., transcribing notes for records, scanning/uploading patient files, etc.), and assisting with patient registration/screening (frontline).
4. Work/rest cycles for prolonged management of potentially overwhelming numbers of critical care patients.

Figure 4. Tiered Staffing Strategy



Modified from the Ontario Health Plan for an Influenza Pandemic Workgroup. Critical Care During a Pandemic.

Case Example: Staffing Approach used by TF MED 14

A tiered approach assisted strategic staff management. Pandemic staffing ratios were used as a guideline, but modified based on tactical patient care/workload. Permitting unit skill sets, nursing personnel were categorized into COVID-19 and trauma teams. A minimum of two staff members were required on shift regardless of census and a critical care nurse for Rapid Response.

Table 3. COVID-ICU

Patient: Nurse Ratio	
Staff	Patient Volume/Acuity
66S Intensive Care RN	2 severe or 10 moderate
66H Medical-Surgical RN	1 severe (with oversight of 66S) or 10 moderate
68C Licensed Practical Nurse	10 moderate

Table 4. COVID-ICW

Patient: Nurse ratio by MOS	
68W Medic	10 moderate (with oversight by 68C or 66H)

RESOURCES:

[Crisis Level Surge – Composition and Roles of the Triage Team, Covid-19 Practice Management Guidelines v7, 3 March 2021, p139](#)

SKILLS BUILDING/TRAINING**Cross-training/Up-training**

All healthcare personnel should be up-trained and practice at the top of their licensure. All healthcare personnel should familiarize themselves with or review how to conduct certain critical activities (in case of personnel shortages):

1. Use of critical care medications and ward stock.
2. Daily preventative maintenance checks and services (PMCS) of critical equipment.
3. Oxygen burn rate for patient usage and refill plan.

The following are recommended critical care topics for providers to review and drill to improve response and success in managing COVID-19 critical patients:

1. Code Blue management of COVID-19 patient.
2. Rapid Sequence Intubation roles of nursing staff.
3. Ventilator management and Arterial Blood Gas (ABG) interpretation.
4. Titration of critical care medications and side effects.
5. PPE donning and doffing.
6. Rapid response team parameters.
7. Proning of a patient in an austere environment.
8. Working knowledge of critical equipment (POGS, EDOCS).
9. Unit specific standard operating procedures.

Units should conduct practice drills: PPE donning/doffing, patient procedures, transfers, proning, Code Blue, etc.

Leadership should consider cross-training non-medical personnel to assist clinical staff as appropriate and allowable. Non-medical personnel must perform [Just-in-Time Training: DHA US-001 HIPAA and Privacy Act Training](#). CAC required.

Society of Critical Care Medicine provides free COVID-19 Resources for Non-ICU Clinicians. Topics include:

- Recognition and Assessment of the Seriously Ill Patient
- Critical Care for the Non-ICU Nurse
- Airway Management
- Airway Assessment and Management
- Diagnosis and Management of Acute Respiratory Failure
- Mechanical Ventilation
- Prone Positioning
- Diagnosis and Management of Shock

RESOURCES:

Crisis Level Surge – Critical Care Triage Tool Example, [DoD COVID-19 Practice Management Guidelines, 3 March 2021. p138](#)

[The Elsevier, COVID-19 Health Care Hub](#) provides clinical toolkits, podcasts, expert opinions and many other tools for providers and staff in the COVID-19 response.

There are also resources for care of older adults and children. The web-based resources are free but require initial completion of a basic online form <https://www.sccm.org/COVID19>

The Defense Institute for Medical Operations (DIMO, www.dimo.af.mil) hosts the following courses:

- MASL D319329 - Medical Readiness Principles of Food Protection
 - MASL D319328 - Malaria & Other Infectious Disease Threats
 - MASL D319318 - Infection Control in Ebola & Pandemic Management
-

STAFF SURVEILLANCE AND STAFFING DECISIONS

Individual risk assessment and fitness for duty for exposed healthcare workers should be determined with the support of COVID testing results, if available, and updated staff medical records. Updated CDC guidelines regarding healthcare workers can be found in the following resources list. In general, the following principles should be followed:

- All healthcare workers should engage in appropriate education, training and policies to comply with infection prevention and control.
- Any healthcare worker who develops fever or symptoms consistent with COVID-19 should immediately self-isolate and contact their established point of contact for testing.
- Per the CDC, “The utility of serologic testing to provide evidence of prior infection that would permit exclusion from quarantine has not been established and is not recommended for this purpose at this time.”
- Healthcare workers who are ill should not be involved in direct patient care.

Within the operational environment, health protection measures must be able to make allowances for mission-critical operational demands beyond the healthcare workers. Certain units, personnel, and functions are mission-critical and incapable of quarantining away from work for 14 days. To overcome this need, certain locations have established the concept of a “firewall” quarantine. While not an official CDC evolution, these firewall quarantines are designed to provide maximal protection from COVID-19 spread while allowing continued operations. It is modeled after the CDC’s recommendation for “Interim U.S. Guidance for Risk Assessment and Work Restrictions for Healthcare Personnel with Potential Exposure to SARS-CoV-2.”

Firewall quarantine is a self-sustaining group of individuals exposed to COVID-19 that must continue to work to sustain mission-critical capabilities. For 14 days, these individuals quarantine and self-monitor for symptoms of COVID-19. They are only allowed in their designated workstation or berthing unit and must wear N95s whenever outside of berthing. Firewalled individuals are required to have separate basic life support (bathrooms, showers) from others who are not firewalled. They also use separate workstations from those who are not quarantined. Meals are delivered, and they are prohibited from visiting the barber, gym, PX, DFAC, etc., until the firewall quarantine completes. Any violations of the firewall quarantine will cause a restart of the 14-day quarantine for the whole firewall cohort and may subject the offender to UCMJ disciplinary actions. Examples of units that have successfully implemented firewall quarantine include healthcare facilities, base defense units, and command and control elements.

RESOURCES:

[CDC Return to Work for Healthcare Personnel with Confirmed or Suspected COVID-19](#)

[Interim U.S. Guidance for Risk Assessment and Work Restrictions for Healthcare Personnel with Potential Exposure to SARS-CoV-2](#)

[Strategies to Mitigate Healthcare Personnel Staffing Shortages](#)

Additional Training, Resources, and Tools are listed at [Appendix B](#)

TESTING

TYPES OF COVID TESTS

There are two broad types of COVID lab tests, distinguished by the method they use to detect infection:

1. Viral Testing – Per CDC, any authorized assay for viral testing that detect either SARS-CoV-2 nucleic acid or antigen. Recommended to diagnose acute infection of both symptomatic and asymptomatic individuals, to guide contact tracing, treatment options, and isolation requirements. There are two major subtypes:
 - a Antigen Testing - fast (<15 minutes) on-site testing that identifies COVID-specific structural components. Relatively inexpensive, but also less sensitive (higher chance of missing positive cases). Previously referred to as “Rapid Antigen Tests.” Renamed by the CDC in December 2020. May require Nucleic Acid Amplification Tests (NAAT) confirmatory testing under certain circumstances. Current FDA EUAs for antigen tests limits their use to symptomatic patients within a certain number of days of symptoms onsets. Special authorization was granted to the DoD by the FDA to use for serially screening asymptomatic individuals in certain circumstances.
 - b NAATs - slower (1 hour - 2 days) in-lab testing that initiates an exponential multiplication of COVID nucleic acid. More expensive, but higher sensitivity (lower chance of missing positive cases) and specificity (lower chance of falsely identifying a true negative as positive) compared to antigen tests. Most common subtype is reverse transcription-polymerase chain reaction (RT-PCR) testing.
2. Antibody Testing - in-lab test identifies the immune response to COVID (antibodies). The FDA has not authorized antibody tests to diagnose SARS-CoV-2. It is not approved for diagnosing current infection but may help determine whether an individual was previously infected. Issues with cross-reactivity to other similar virus antibodies. Commonly used for surveillance testing and epidemiologic studies.

RESOURCES:

[Overview of Testing for SARS-CoV-2 \(COVID-19\)](#)

[Interim Guidance for Antigen Testing for SARS-CoV-2](#)

[Using Antibody Tests for COVID-19](#)

COVID TESTING STRATEGIES

There are also three broad categories for COVID testing strategies by the CDC:

1. Diagnostic - used to identify current infection in symptomatic patients or asymptomatic close-contacts.
2. Screening - used to identify infected persons who are asymptomatic and without known or suspected exposure. Screening testing is performed to identify persons who may be contagious so that measures can be taken to prevent further transmission.
3. Surveillance - used to gain information at a population level with de-identified results returning in aggregate and not to the individual. It is intended to monitor community - or population-level outbreak of disease, or to characterize the incidence and prevalence of disease.
4. Surveillance testing cannot be used for individual decision-making.

Table 5. CDC COVID TESTING STRATEGIES

CDC Summary of Testing Strategies for SARS-CoV-2			
	Diagnostic	Screening	Surveillance
Symptomatic or known or suspected exposure	Yes	No	N/A
Asymptomatic without known or suspected exposure	No	Yes	N/A
Characterize incidence and prevalence in the community	N/A	N/A	Yes
Results may be returned to individuals	Yes	Yes	No
Results returned in aggregate to requesting institution	No	No	Yes
Testing can be performed in a CLIA-certified laboratory	Yes	Yes	Yes
Testing can be performed in a non-CLIA-certified laboratory	No	No	Yes
Test system must be FDA-authorized or be offered under the policies in FDA's guidance	Yes	Yes	No

Further details are available at the CDC website under "Testing Overview" in the Healthcare Workers section.

RESOURCES:

[Overview of Testing for SARS-CoV-2 \(COVID-19\)](#)

[Frequently Asked Questions about Coronavirus \(COVID-19\) for Laboratories](#)

SAMPLE POOLING

When disease prevalence in an area is sufficiently low, certain Nucleic Acid Amplification Testing (NAAT) diagnostic testing systems are approved for sample pooling in order to save time and Class VIII materials. However, the combination of multiple specimens into one test dilutes the samples resulting in lower sensitivity. Theater lab officers should establish area-specific guidelines for pooling thresholds based on disease prevalence and specific equipment limitations. For example, the BioFire EUA (see the following resources list) includes three pages on approved pooling techniques. Lastly, pooling should only be executed at CLIP-certified laboratories.

The two most common NAAT systems in CENTCOM include the BioFire Diagnostic system and the GeneXpert family of diagnostic systems. At the time of this writing, the BioFire is approved for pooling samples; the GeneXpert systems are not approved for pooling. Pooled testing can be used for screening or diagnostic testing.

RESOURCES:

[BioFire® COVID-19 Test Instructions for Use](#)

[Interim Guidance for Use of Pooling Procedures in SARS-CoV-2 Diagnostic, Screening, and Surveillance Testing](#)

DoD TESTING ALGORITHMS

Refer to current [DoD Corona Virus Task Force COVID-19 Testing Protocols](#)

MEDICAL MANAGEMENT

IMMEDIATE CARE

There is no cure for COVID-19. The treatment available now remains supportive care, which includes antipyretics, supplemental oxygen, and potentially mechanical ventilation. Medications currently used in the treatment of COVID-19 are off-label, are a part of research study protocols, or have been released by the FDA for Emergency Use Authorization.

COVID-19 has manifested primarily as a significant respiratory infection that can rapidly deteriorate into a severe pneumonitis, requiring supplemental oxygen and potentially mechanical ventilation. The severity seems to primarily affect the elderly population, and those with underlying medical comorbidities. However, this virus has also affected otherwise young healthy individuals. Providers must be aware and monitor all patients for potential rapid deterioration and intervene as soon as possible.

The DoD developed the [DoD COVID-19 Practice Management Guidelines](#) to assist providers throughout the DoD with comprehensive care for treatment and management of COVID-19.

With the decreased availability of staff and resources in the austere setting, a supplement to the DoD PMG for COVID-19 contains a section for considerations in the austere environment, under the same link immediately above.

PROLONGED CARE

Due to the global nature of this pandemic, as well as the infectivity of COVID-19, there are many special considerations when planning to evacuate patients from downrange. It is recognized that the DoD's usual protocols and ability to evacuate patients has been severely limited. Patients may require treatment over a prolonged time (possibly several days) prior to evacuation to a higher echelon of care. It is a priority for the downrange provider to plan for longer than expected care. The DoD PMG for COVID-19 and the CPG for austere locations referenced previously, are useful tools. Additionally, the [JTS prolonged field care guidelines](#), while not specific to COVID-19, are useful for prolonged field care and can be found within the following resources links.

RESOURCES:

[JTS Nursing Interventions in Prolonged Field Care CPG, 22 Jul 2018](#)

[JTS Documentation in Prolonged Field Care CPG, 13 Nov 2018](#)

DISCHARGE

Discharging a patient from transmission-based precautions (TBP) AKA isolation should follow the CDC symptom based strategy. Patients can move through the various levels of care (mild outpatient self-care, moderate ward level care and severe ICU level care) but must remain in isolation and follow TBPs during this entire time.

For most adult patients with COVID-19 illness, isolation and precautions can be discontinued 10 days after symptom onset AND after resolution of fever for at least 24 hours, without the use of fever-reducing medication, AND with improvement of other symptoms.

Some adults with severe illness may produce replication-competent virus beyond 10 days that may warrant extending duration of isolation and precautions for up to 20 days after symptom onset.

For adults who never develop symptoms, isolation and other precautions can be discontinued 10 days after the date of their first positive RT-PCR test result.

RESOURCES:

[Interim Guidance on Duration of Isolation and Precautions for Adults with COVID-19](#)

TELEMEDICINE

Listed below are emergency telehealth resources which forward-stationed medical teams/medics can use for assistance or clarification on any topics or concerns:

VITAL-T Program

Virtual Inspection and LINKUP in Theater (VITAL-T) provide “real time” virtual access to Quality and Safety (QS) expertise through Virtual Health (VH) capabilities for Infection Prevention and Control (IPC), Patient Safety (PS), and Medication Safety (MS).

Submit a Vital-T Consult in 2 Easy Steps

1. Call with questions or to arrange a Virtual Consult or Service with the appropriate QS expert.
2. Email: <mailto:usarmy.jbsa.medcom.list.medcom-vitalt@mail.mil> or
24-hour VITAL-T hotline: 210-307-0923

During the COVID-19 crisis, consult services could be conducted using any of the following approved means (the optimal means for connection will be individualized):

1. Google Duo
2. FaceTime
3. Skype
4. Adobe Connect

AD.VI.S.OR

[Advanced Virtual Support for Operational Forces](#) (AD.VI.S.OR) program is specifically designed for operational virtual health support. Phone: 833-238-7756; Email: dod.advisor_office@mail.mil

Additionally, many Virtual Critical Care Consultation (VC3) service providers have deployed to austere settings before and can help work through the unique problems faced in austere settings.

DHA Infection Prevention and Control Tiger Team

The DHA Tiger Team is an alternative source for infection prevention and control reach-back, which is overwhelmed in the current situation. The team will review received questions on a daily basis and work to develop a response within 1-2 business days dha.ncr.clinic-support.list.ipc-group@mail.mil.

Table 6. Local USCENTCOM ADVISORS for Urgent Needs

<i>Local USCENTCOM ADVISORS for Urgent Needs</i>
379 EMDG SGH: DSN 318-455-5042 or cell 011-974-5080-3442
379 EMDG Help Line (answered 24/7): DSN 318-455-1000
455 EMDG: DSN 318-481-4664
Kabul: DSN 318-449-9259
BDSC: DSN 318-239-2033
USMH-K: 318.430.2508
USCENTCOM Preventive Medicine Physicians: DSN 312-529-0348
USCENTCOM Pharmacist: DSN: 318-480-5094/SVOIP: 308-430-6834
PHEO: DSN is 318-480-6087; SOVIP is 308-430-8507

As a final alternative, the following MEDCENS below may be contacted (ask for the on-call critical care staff).

****Note:** MEDCENS may be intensively involved in COVID-19 response locally. Personnel should exhaust all local, regional, USCENTCOM reach back and all dedicated telehealth resources PRIOR to calling MEDCENS.

Table 7. MEDCEN Contact List

<i>MEDCEN Contact List</i>
Landstuhl Regional Medical Center, Germany. DSN: 314-590-7141 Intensive Care Unit
Walter Reed National Military Medical Center, MD. (301) 295-4611, option 4 Command Duty (301) 295-4810 Emergency Room
Madigan Army Medical Center, Fort Lewis, WA (DSN 782) (253) 968-1110 Information Desk
Brooke Army Medical Center, Fort Sam Houston, TX (DSN 429) (210) 916-0808 Emergency Room
Naval Medical Center Portsmouth, NS Norfolk, VA (DSN 377) (757) 592-5473 Critical Care (757) 953-1365 Emergency Room
Eisenhower Army Medical Center, Fort Gordon, GA (DSN 773) (706) 787-6938/6019 AOD (706) 787-6039 Emergency Room
David Grant Medical Center, Travis Air Force Base, CA (DSN 799) (707) 423-3040 ICU or (707) 423-3825 Emergency Room
Tripler Army Medical Center, HI (DSN 433) (808) 433-6661 Information Desk (808) 433-4032 ICU or (808) 433-3707 Emergency Room
William Beaumont Army Medical Center, Fort Bliss, TX (915)892-6880 House Supervisor or (915) 742-2139 ICU
Keesler Medical Center, Keesler AFB, MS (DSN 591) (228) 376-0500 Emergency Room

BEHAVIORAL HEALTH

Leaders **must** consider the behavioral health and mental/physical resiliency of health providers and staff responding to an outbreak, as well as that of COVID-19 patients and the general population, who are experiencing major life-style changes and isolation. A number of tools and resources have been provided for leaders to respond to behavioral health concerns and improve resiliency. Forward stationed medics/medical teams are encouraged to reach back locally to Role 2/Role 3 facilities, where behavioral health resources may be stationed, utilize chaplain services, and telehealth resources to connect persons in need with help.

Special considerations for behavioral health during the COVID-19 pandemic include:

1. Prolonged isolation of COVID-19 designated staff and patients.
2. Disruption of non-duty activities and resiliency behaviors (e.g., physical activity, rest/relaxation).
3. Risk for minimal work/rest cycles due to potentially overwhelming numbers.
4. Moral stressors of triaging patients to receive care in the pandemic environment.
5. Fear response for novel or unfamiliar risks/threats.

RESOURCES:

Behavioral Health and Wellness in Covid-19 Clinical Management (patients, providers) - [DoD COVID-19 PMG v7.0, 3 March 2021 p93-96](#).

[Resiliency/Well-Being – Stress and Coping, CDC Mental Health and Resiliency during COVID-19](#)

Leadership checklist to mitigate team stress is located within [Appendix C](#).

Leadership checklist to promote team sleep is located within [Appendix D](#).

[Navy Leader’s Guide for Managing Sailors in Distress](#) – The purpose is to help Leaders recognize distress related behaviors, provide support to Sailors within the unit, and collaborate with Navy helping agencies to meet the needs of distressed individuals.

The DoD and VA offer the below mobile apps for mental health guidance.

Provider Resilience: Through psychoeducation and self-assessments, Provider Resilience gives frontline providers tools to keep themselves productive and emotionally healthy as they help our nation’s service members, veterans, and their families [Apple](#) and [Android versions](#).

Psychological First Aid (PFA): PFA Mobile was designed to assist responders who provide psychological first aid (PFA) to adults, families, and children [Apple](#) and [Android versions](#).

Other apps can be viewed by downloading the below brochure at <https://health.mil/Reference-Center/Publications/2019/08/28/DoD-and-VA-Mobile-App-Clinicians-Guide>. The Brochure does not provide links to the apps.



Figure 5. Clinician's Guide of Mobile Health Apps

CLINICIAN'S
GUIDE

Department of Defense and Veterans Affairs
Mobile Health Apps for *PATIENTS*

Presenting Conditions

App Features

	ACT Coach	AIMS	Army OneSource Money Matters	Army PRT	The Big Moving Adventure	Breathe2Relax	CBFI Coach	Concussion Coach	CPT Coach	CSF2 Goal Setting	Dream EZ	High Intensity Tactical Training	LifeArmor	Mindfulness Coach	Mood Coach	MOVE! Coach	Moving Forward	MyPlate	Parenting2Go	PE Coach 2	Performance TRIAD	Physical Readiness Training	Positive Activity Jackpot	PTSD Coach	Sesame Street for Military Families	STAIR Coach	Stay Out Coach	T2 Mood Tracker	Tactical Breather	VetChange	Virtual Hope Box
Alcohol/Drugs/Tobacco																															
Anger/Irritability																															
Anxiety/Stress/Depression																															
Family/Social																															
Headaches/Pain																															
Mindfulness																															
Nutrition/Exercise																															
Personal development/Goal setting																															
Post Traumatic Stress																															
Resilience																															
Sleep																															
Spirituality																															
Traumatic Brain Injury																															
																															
																															
Corresponding Guide or Handout																															
Password Protected																															
To Be Used with Manualized Treatment																															

ARMED FORCES MEDICAL EXAMINER SYSTEM AND MORTUARY AFFAIRS

All U.S. Military Service Members and U.S. Civilian personnel who die in an Operational Environment are under the medico-legal jurisdiction of Armed Forces Medical Examiner System (AFMES) in accordance with Title 10 U.S. Code 1471. If a death occurs, immediately notify AFMES at (202) 409-6811. All medical intervention should remain in place. Contact mortuary affairs for further guidance on the handling of the decedent. Secure clothing, personal protective gear, and medical/treatment records. These should accompany the decedent to mortuary affairs.

RESOURCE:

[Safety and Health Guidance for Mortuary Affairs Operations: Infectious Materials Technical Guide 195A, November 2015.](#)

ANCILLARY SERVICES

Ancillary service personnel will practice previously described infection prevention and COVID-19 medical management control measures.

PHARMACY

Goals are to reduce the impact of exposure risks and preserve access to medications. Pharmacists will support ongoing clinical evaluation studies and treatment protocols for emerging therapeutics.

RADIOLOGY

Once providers deem a diagnostic study necessary, take measures to minimize risk of cross contamination of equipment and environment. Equipment (including wheels) and X-ray cassettes shall be wiped down prior to entering and exiting patient care areas. Use disposable X-ray cassette protective sleeves or other type of similar barrier material for portable chest X-ray capability and equipment. Portable machines should be positioned so as to prevent contact with the patient whenever possible.

LABORATORY (SCREENING/TESTING)

Appropriate PPE and infection control/prevention procedures to prevent blood borne pathogen and aerosol exposure will be employed. Patient screening may include nasopharyngeal swab specimens (most common) or tracheal aspirates from intubated patients.

RESOURCES:

[Nasopharyngeal swab technique training video \(7:19\)](#)

[Management of COVID-19 in Austere Operational Environments v3.0, 23 Apr 2021, p9-10](#)

[DoD COVID-19 PMG v7.0, 3 March 2021, p22](#)

[Frequently Asked Questions about Biosafety and COVID-19](#)

ENVIRONMENTAL SERVICES/HOUSEKEEPING

Employed cleaning protocols should ensure adequate sanitization in all environments, including quarantine/isolation/patient care areas, as well as all workspaces and quarters. All non-dedicated, non-disposable medical equipment used for patient care should be cleaned and disinfected according to manufacturer's instructions and facility policies.

- *Routine Cleaning and Disinfection Procedures:* Use cleaners and water to pre-clean surfaces prior to applying an EPA-registered, hospital-grade disinfectant to frequently touched surfaces or objects for appropriate contact times as indicated on the product's label.
- *High Touch Surfaces:* In addition to standard environmental cleaning, employ routine cleaning of high-touch surfaces: tables, chair arm rests, doorknobs, light switches, countertops, handles, desks, phones, keyboards, mouse devices, toilets, faucets, sinks, etc.
- *Bathroom Entrances/Exits:* When possible, position trash cans inside bathrooms near the door to allow a no-touch exit. Hand washing signs should be placed in entryways.

RESOURCES:

[List of Disinfectants for Use against SARS-CoV-2](#)

[Interim Infection Prevention and Control Recommendations for Patients with Suspected or Confirmed COVID-19 in Healthcare Settings, 10. Implement Environmental Infection Control](#)

FORCE HEALTH PROTECTION

Force Health Protection (FHP) assets which include the public health emergency officer (PHEO), public health, preventive medicine, and veterinary services are important stakeholders and subject matter experts on issues of infectious disease control (including zoonotic disease), sanitation, food protection and safety, and water systems protection and safety. Veterinary Services (VS) are equally important subject matter experts in animal medicine and husbandry, human-animal bond, and agricultural systems. Where available, FHP personnel should be consulted regarding these issues; there are now extensive Force Health Protection (FHP) Guidance documents published by the DoD available here: <https://www.defense.gov/Explore/Spotlight/Coronavirus/Latest-DOD-Guidance/>

A series of continually updated supplemental guidance documents outline the DoD position on all aspects of force health protection related to COVID-19 to include: testing, restriction of movement, diagnosis, recovery, travel restrictions, patient movement, non-pharmaceutical interventions (mask wear, social distancing, building occupancy restrictions, etc.), return to work criteria, and health protection conditions (HPCON). These FHP supplements are updated to include the evolving understanding about COVID-19 and changes in CDC and FDA recommendations and regulations. USCENTCOM policy is closely aligned with the DoD FHP documents but health care personnel must be familiar with both and should look to the link above regularly for updated information.

This information in this section is intended to supplement not supersede the USCENTCOM/Component and TRANSCOM policy and guidance. Any operational issues or challenges regarding FHP or the movement and care of authorized animals and animal-care personnel should be resolved through local, ARCENT, and USCENTCOM Command leadership.

[Contact tracing](#) is a critical part of COVID-19 mitigation, and must be supported by the command group. COVID Trace Team members will be required to complete training (see [Appendix B: Additional Training and Resources](#)). All positive cases will be [contact](#) traced and all individuals identified as a close contact will be placed in quarantine based on current USCENTCOM policy and DoD FHP guidance supplements. All confirmed cases will be [contact](#) traced and identified; close contacts will be placed in quarantine. Release from quarantine and isolation should follow current DoD and USCENTCOM policy but local commanders may impose more restrictive measures.

FIELD SANITATION AND ISOLATION

In many cases, isolation facilities will need to be rapidly established that may require alternative waste and hygiene approaches to protect the uninfected general population (e.g. alternative showering facilities, pit latrines, bed cans). FHP assets should be consulted to ensure best possible practices are being implemented and that all safety and handling precautions are implemented.

RESOURCE:

[Management of COVID-19 in Austere Operational Environments v3.0, 23 Apr 2021](#)

[Central Command Regulation 40-2 08 Nov 2017, Deployment Force Health Protection](#) (SIPRNet)

[CDC Sanitation & Hygiene](#)

[DoDD 6200.04, Force Health Protection, Oct 2004](#)

[ATP 4-02.8, Force Health Protection, Mar 2016](#)

[Training Circular 4-02.3, Field Hygiene and Sanitation](#)

[Health Service Support Casualty Prevention for Expeditionary Operations, AFTTP 3-42.2, 2004](#)

FOOD SAFETY AND FOOD SERVICE

There is still no evidence for concern that food and food packaging serves any role in transmission of COVID-19. Protection in dining facilities will focus on protection against direct human-to-human transmission and should include the following considerations.

- A la carte dining options (e.g., deli and sandwich stations, omelet stations) should be suspended during the pandemic.
- Fresh fruits and vegetables and deserts should not be presented in a salad bar format. These foods should be cleanly prepared and packaged in covered food containers.
- Meal lines and dining areas should consider social distancing. Seating should be placed in appropriate distances and/or limitations on capacity should be implemented. Consider outdoor dining options and take out approaches.
- Consider implementing servers – designated personnel that are screened for fever and symptoms at the start of each shift - instead of self-serve approaches or pre-plating or packaging meals to reduce contact during the serving process.

RESOURCE:

[DoD Veterinary Services COVID-19 Resource Page](#) Provides white papers, external links, situation reports and other operational guidance related to food safety and protection, FHP, and animal medicine/animal health during the COVID-19 pandemic.

[FDA Best Practices for Retail Food Stores, Restaurants, and Food Pick-Up/Delivery Services During the COVID-19 Pandemic](#)

VETERINARY CARE

Government owned animals (GOAs) and authorized privately owned animals (POAs) should continue to receive veterinary medical care in accordance with DoD and contact specific guidelines at local VS locations (veterinary treatment facility or similar). Any currently understood possibility of zoonotic transmission DOES NOT exceed the harm caused by not ensuring care for these animals. Elective procedures and veterinary medical care activities should be cancelled for the duration, but standards of care for public health and readiness should be maintained in accordance with applicable policy, health certificate requirements, and best practice guidelines (e.g. vaccination, emergency care, retail/distribution of medication and preventatives, etc.).

RESOURCES

[DoD Veterinary Services COVID-19 Resource Page](#)

[CDC COVID-19 and Animals](#)

[American Veterinary Medical Association \(AVMA\) COVID-19 webpage](#)

ANIMAL TO HUMAN TRANSMISSION

The COVID-19 virus is known to have zoonotic origins, believed to originate at a live animal and wet market in Wuhan, China. It appears that in rare situations, COVID-19 infected persons can spread the virus to animals. The virus has been detected in felids (domestic cats, lions, tigers, snow leopards, and cougar), canids (domestic dogs), non-human primates (gorillas), and mustelids (ferrets, farmed mink and wild mink). It is unlikely that this virus will cause significant clinical disease in healthy domestic animals.

There is no evidence to suggest domestic animals serve any significant epidemiological role in transmission of the virus to humans. Human-to-human transmission remains the primary source of transmission. However, given the possibility for zoonotic transmission of the COVID-19 virus, the following recommendations should be implemented:

- Commanders should strictly enforce USCENTCOM General Order (GO) 1C (paragraph 2.g) that prohibits the adoption of unauthorized pets or mascots of any kind on DoD installations. Exceptions to this policy include authorized GOAs, including military working dogs (MWDs) and DoD Contracted Working Dogs, and installations where personnel live off-base and are authorized POAs.
- ***It is not recommended to routinely test animals for COVID-19*** using BioFire or any other testing platform. The CDC provides guidance on testing animals for COVID-19. The CCSG team and CENTCOM Command Veterinarian should be consulted PRIOR to any testing of animal patients for COVID-19.

RESOURCES

[USCENTCOM General Order 1C \(para. 2g\), 21 May 2013](#)

[AVMA Interim recommendations for intake of companion animals from households where humans with COVID-19 are present](#)

[NASPHV Compendium of Measures to Prevent Disease Associated with Animals in Public Settings](#)

[Interim Infection Prevention and Control Guidance for Veterinary Clinics Treating Companion Animals During the COVID-19 Response](#)

[CDC Evaluation for SARS CoV-2 Testing in Animals](#)

[DoD Guidance for Coronavirus Disease 2019 and Animals](#)

CONSIDERATIONS FOR ANIMAL CARE PERSONNEL

VS personnel should routinely implement and train local non-VS animal care personnel in the principles and standards of infection control outlined in the [NASPHV Compendium of Measures to Prevent Disease Associated with Animals in Public Settings](#) and applicable infection control policy and best practices.

Elective veterinary visits and procedures should be cancelled through the pandemic. Animal care personnel should remain aware of current shortages and high demand of PPE for their human medical counterparts and practice prudent PPE implementation when treating animals, conducting laboratory procedures, and routine husbandry practices during the COVID-19 pandemic. The CDC makes recommendation on their website for PPE use when caring for animals. Working dog handlers that

require isolation or quarantine should NOT conduct daily care for their dog. The Kennel Master, trainer, or other handlers should provide routine care for that dog until the assigned handler can return to duty.

RESOURCES:

[CDC Interim Guidance for Public Health Professionals Managing People With COVID-19 in Home Care and Isolation Who Have Pets or Other Animals](#)

[NASPHV Compendium of Measures to Prevent Disease Associated with Animals in Public Settings](#)

SUPPLIES

All disposable and non-durable medical supplies (including PPE) stock will be distributed and stocked IAW with pre-planning and role/location specific guidance.

Employ cleaning protocols to ensure adequate sanitization in all environments, including quarantine/isolation/ patient care areas, workspaces and quarters. All non-dedicated, non-disposable medical equipment used for patient care should be cleaned and disinfected according to existing location and USCENTCOM specific infection control policies, manufacturer guidelines, and best practices.

Figure 6. PPE Recommendations for the MHS

Category	Definition	Required Isolation/PPE
0	Patient not suspected of having COVID-19	STAFF <ul style="list-style-type: none"> Surgical mask PPE according to task. See standard precautions. PATIENTS: Avoid close contact with any staff unless necessary for healthcare purposes.
1	Asymptomatic patient with known exposure to COVID-19 OR traveled from high risk areas within last 14 days.	STAFF <ul style="list-style-type: none"> Surgical mask PPE according to task. See standard precautions. PATIENTS: MUST wear surgical mask if traveling outside room for medically essential purposes.
2	Patient under investigation (PUI) or positive COVID-19	STAFF <ul style="list-style-type: none"> Contact precautions (gown and gloves) Droplet precautions (surgical mask appropriate per current CDC guidelines if no aerosol-generating procedures performed in room) Eye protection (face shield or goggles) PATIENTS: MUST wear surgical mask if traveling outside room for medically essential purposes.
3	Positive COVID-19 requiring aerosol-generating procedures (i.e. BIPAP, CPAP, endotracheal intubation, high flow nasal cannula, nebulizers, tracheal suctioning)	STAFF <ul style="list-style-type: none"> Contact precautions (gown and gloves) Consider head and foot covers Airborne precautions (N95 respirator or PAPR) Eye protection (face shield or goggles) Negative pressure room PATIENTS: MUST wear surgical mask if traveling outside room for medically essential purposes.

Source: Figure 8: PPE Recommendations for the MHS (adopted for the MHS using CDC guidelines accessed 9 Jun 2020; <https://www.cdc.gov/coronavirus/2019-ncov/hcp/using-ppe.html>) PAPR: Powered air-purifying respirator. PUI: Patient under investigation. High-risk area- Area with level 3 travel health notice identified by the CDC.

RESOURCES:

[CDC Strategies to Optimize the Supply of PPE and Equipment](#)

[HHS Hospital Personal Protective Equipment Planning Tool](#)

[Personal Protective Equipment \(PPE\) Burn Rate Calculator](#)

POINTS OF CONTACT

The USCENTCOM Command Surgeon (CCSG) team is the main point of contact for this document at (813)-529-0345/0361/0362 (COMM/DSN) or centcom.macdill.centcom-hq.mbx.ccs-g-clinops@mail.mil

LTC Fred Hauser and COL Vinh Tran are the USCENTCOM Public Health Emergency Office point of contacts and can be reached through the same organization mailbox (centcom.macdill.centcom-hq.mbx.ccs-g-clinops@mail.mil)

INDEX

AE Patient Classification	1
Airborne Spread	1
Ancillary Services	29
Antibody/Antigen	2
Behavioral Health	27
Cleaning, Disinfecting Households and Facilities	9
Contact Tracing	38
Critical Resource Conservation	12
Command and Control	4
Communications	5
Contact Spread	2
Crisis Standards of Care	3
Definitions	1
Documentation	6
Droplet Spread	2
Environmental Services/Housekeeping	29
Executive Summary	1
Facilities Management	8
Field Sanitation and Isolation	31
Ethical Considerations	13
Human Resources	18
Hygiene, Social Distancing, Shelter in Place, and General Protection Measures	17
Infection Prevention and Control (IPC)	25
Intermediate Care Ward (ICW)	2
Laboratory (Screening/Testing)	29
Medical Centers (MEDCENS), Contact Information	26
Medical Management	24
Mortuary Affairs	28

Operational Planning	4
Pathogen	29
Patient Under Investigation (PUI)	2
Performance Improvement	6
Personal Protective Equipment (PPE) Considerations	8
Pharmacy	29
Quarantine Facilities	10
Radiology	29
Resiliency/Well-Being	27
Screening/Testing	23
Skills Building/Training	19
Staff (Patient: Staffing Ratios, Resilience/Well-Being)	18
Staff Surveillance and Staffing Decisions	21
Supplies	33
Surge Capabilities	11
Suspected, Confirmed Cases	6
Telemedicine/VITAL-T/Points of Contact	25
Transport	7
Triage/Security	14

APPENDIX A: PANDEMIC RESPONSE PLANNING CHECKLIST

Role of Care: • R1 • ERC • R2 • R3 • R4
 Primary POC Email: Phone:

1. Pandemic Threat Working Group Members, Principle Command & Control, Medical Lead		
Identify Members of Threat Working Group <ul style="list-style-type: none"> Medical Lead Recommended: Infectious disease specialist or emergency medicine/surgeon, if medical specialist unavailable. Minimal Recommended: BOS-I Commander or designee, Medical Asset Leads or Designated Specialist, Public Affairs, Medical Assets, Tenant Unit/Mission Representatives, Security Forces. 		
	Quantify threat to force	
	Plan for dissemination of information to MTF population	
	Determine Health Protection Condition (HPCON)	
	Identify primary sources of threats and screening methods (i.e. front gate, flight line)	
2. Centralized Concerning Symptom Reporting		
	Establish diagnosis algorithm for concerning disease	
	Establish hotline to call in symptoms to arrange meetup between protected medical asset and possible infected patient	
	Designate evaluation area segregated from healthy population	
	Closure of medical assets to unscreened patients	
3. Designated healthy, quarantine, and Infected/Isolated Facilities		
	Plan social separation: separate population into healthy, quarantine, and infected/isolated populations	
	Establish means to provide sleeping, food, water, hygiene, sanitation and socialization/recreation	
	Method to enforce segregation of population	
	Establish criteria for entry and exit of patients from groups (i.e. clinically resolve or infected) and rules for outside personnel entry (i.e. medical providers only with specific PPE)	
	Establish decontamination/disinfection protocols	
4. Treatment Plan		
	Identify locations that can provide medical support (ward/ICU) and maintain isolation	
	Designation of treatment team and contingencies for medical team casualties	
	Inventory personal protection equipment, materials, and medications and report to higher command estimated duration of supplies	
	Request any additional assets from higher commands that may be required	
	Biohazard disposal plan	

5. Specific Equipment		
	Respirators with filter capacity of at least 94% - N95 or FFP2 Respirators	
	Disposable gowns, face shields, soap	
	Disinfectant (i.e. bleach, CaviWipes™)	
	Oxygen concentrators for nasal cannula	
	Oxygen tanks with refill capacity or oxygen source (i.e. POGS) capable of providing pressurized oxygen to enable FiO ₂ of 100%	
	i-STAT with arterial blood gas cartridges	
	Thermometer	
	Tracheal suction catheters and suction (inline)	
	Bag valve masks	
	Pulse oximeter	
	Ventilator with air inlet filters	
	Ventilator circuits	
	Heat and moisture exchanger (HME) and bio expiratory filters for ventilator circuits	
	Nasogastric tubes	
	Long term sedatives (i.e. Propofol, Ketamine) and paralytics (i.e. Vecuronium, Pancuronium)	
	Deep vein thrombosis prophylaxis (Chemical and Mechanical)	
	Antibiotics and supportive care medications dependent on prevalent pathogens	
6. Evacuation plan		
	Establish evacuation protocols to higher levels of care versus treat in place	
	Verify function and availability of in theater patient movement capabilities	
	Plan for disposition of any contagious human remains	
7. Contingencies for contractors/civilians		
	Plan for chronic medication shortages and medical care as mail, movement, and local facilities become unavailable	

APPENDIX B: ADDITIONAL TRAINING AND RESOURCES


The following resources are available for rapid access and review in preparing for or responding to COVID-19 patients locally:

1. *Current COVID-19 Resources: Policy, Military-specific, Clinical*, etc. will be posted in the DoD COVID-19 Clinical Operations Group site
<https://www.milsuite.mil/book/groups/covid-19-clinical-operations-group>
2. *DoD Instruction for Public Health Emergency Management (PHEM) within the DoD*. DoD6200.03 (28 Mar 2019)
<https://www.esd.whs.mil/Portals/54/Documents/DD/issuances/dodi/620003p.pdf>
3. *Joint Trauma System (JTS) Clinical Practice Guidelines (CPGs)*
https://jts.amedd.army.mil/index.cfm/PI_CPGs/cpgs
4. *The Elsevier, COVID-19 Health Care Hub*. provides clinical toolkits, podcasts, expert opinions and many other tools for providers and staff in the COVID-19 response.
<https://elsevier.health/en-US/covid-19/home>
5. *The Elsevier Engineering Resources* for the COVID-19 response
<https://www.elsevier.com/connect/engineering-resources-for-the-covid-19-response>
6. *Infection Prevention and Control Policy in a Deployed Setting* and USCENTCOM Surgeon (CCSG) Clinical Operations SharePoint site (CAC Required)
<https://intelshare.intelink.gov/my.policy>
7. *The US DoD Coronavirus Rumor Control* should be used to clarify points of confusion and prevent decision making based on misinformation
<https://www.defense.gov/Explore/Spotlight/Coronavirus/Rumor-Control/fbclid/I/>
8. *The Centers for Disease Control and Prevention*, COVID-19 webpage
<https://www.cdc.gov/coronavirus/2019-ncov/index.html>
9. *National Institute of Health, COVID19 webpage*
<https://www.covid19treatmentguidelines.nih.gov/>
10. *COVID-19: How to Conduct Contact Tracing*, Health Information Products e-Catalog, Army Public Health Center
<https://ephc.amedd.army.mil/HIPECatalog/viewItem.aspx?id=1841>
11. *COVID-19: Contact Tracing Toolkit*, Health Information Products e-Catalog, Army Public Health Center
<https://phc.amedd.army.mil/topics/campaigns/covid19/Pages/Guidance-and-Alerts.aspx>
12. *COVID-19 | Sustaining Operations: Roadmap to Coronavirus Disease Risk Reduction in Army Training Formations*, COVID-19 | SUSTAINING OPERATIONS: Roadmap to Coronavirus Disease Risk Reduction in Army Training Formations, Health Information Products e-Catalog Army Public Health Center
<https://ephc.amedd.army.mil/HIPECatalog/viewItem.aspx?id=1834>

APPENDIX C: MITIGATING TEAM STRESS

 <h1>COVID-19 Leadership Checklist</h1> <h2>Mitigating Team Stress</h2>	
<h3>SHARE INFORMATION</h3> <p>Sharing information establishes communication and trust with your team.</p> <ul style="list-style-type: none"> <input type="checkbox"/> Stay up-to-date on the latest developments <input type="checkbox"/> Share what you know with the team <input type="checkbox"/> Let them know when you don't know the answer <p>Self Check: Have you updated your team recently?</p>	<h3>MAINTAIN PSYCHOLOGICAL RESILIENCE</h3> <p>Using mental resilience skills can help people manage stress and stay strong.</p> <ul style="list-style-type: none"> <input type="checkbox"/> Encourage a balanced diet of news to avoid feeling overwhelmed <input type="checkbox"/> Keep a detailed to-do list to keep things manageable <input type="checkbox"/> Use positive self-talk or buddy talk to get through stressful moments <input type="checkbox"/> Use "Grounding" (name 3 things you can see, hear, and physically feel) to reduce anxiety spikes and orient yourself to the moment <p>Self Check: What mental resilience skills are you practicing?</p>
<h3>CONNECT</h3> <p>Connecting with others can help prevent people from feeling isolated and alone.</p> <ul style="list-style-type: none"> <input type="checkbox"/> Run regular meetings to provide structure and stability <input type="checkbox"/> Strengthen your team's sense of community and shared purpose <input type="checkbox"/> Set up a group text to check in regularly with all team members including those without government iPhones <p>Self Check: Are you connecting with your own leaders and teammates?</p>	<h3>NORMALIZE STRESS</h3> <p>It is important to acknowledge the impact of stress, letting unit members feel more connected and less emotionally isolated.</p> <ul style="list-style-type: none"> <input type="checkbox"/> Recognize your team's stress ("This is uncharted territory") <input type="checkbox"/> Remember there are individual differences in how people cope with stress <input type="checkbox"/> Give permission to talk about stress to the team <input type="checkbox"/> Recognize that high-achievers are likely to feel even more stress during crises <p>Self Check: Have you acknowledged your own stress level to someone?</p>
<h3>RECOGNIZE LIMITS</h3> <p>Stress can diminish people's ability to process complex information.</p> <ul style="list-style-type: none"> <input type="checkbox"/> Remember to repeat whatever is important and over-communicate <input type="checkbox"/> Be patient if someone makes a mistake or isn't tracking <input type="checkbox"/> Build in redundant checks for critical pathways to reduce errors <p>Self Check: Are you making simple mistakes? Do you need to take a minute to recharge?</p>	<h3>SEIZE THE MOMENT</h3> <p>Leaders can reframe this moment as a critical opportunity for the entire team to contribute to the shared mission of finding solutions to the crisis.</p> <ul style="list-style-type: none"> <input type="checkbox"/> Remind your team of the important mission at hand <input type="checkbox"/> Everyone has an essential role to play, no matter their rank or occupation <p>Self Check: How can this challenge provide you a leadership opportunity?</p>
<h3>MAINTAIN PHYSICAL RESILIENCE</h3> <p>When people take care of themselves physically, they can handle stress better.</p> <ul style="list-style-type: none"> <input type="checkbox"/> Prioritize sleep <input type="checkbox"/> Encourage good nutrition <input type="checkbox"/> Get regular exercise <p>Self Check: Are you remembering to take care of your physical health?</p>	<h3>CONTROL THE CONTROLLABLES</h3> <p>Reduce stress and save energy by focusing efforts on what can be controlled and accepting what can't be controlled.</p> <ul style="list-style-type: none"> <input type="checkbox"/> Encourage your team members to identify what they can control <input type="checkbox"/> Have team members practice deep breathing and mindfulness when things start to feel like they are out of control <p>Self Check: What is within your control? What do you have to accept?</p> <div data-bbox="824 1402 922 1541">  </div> <h3>TAKE THE LONG VIEW</h3> <p>This isn't a one-time process. Pace yourself and your team for a marathon. Remember to be kind to yourself and your team.</p> <div data-bbox="1328 1423 1442 1528"> <p>RESEARCH TRANSITION OFFICE CENTER FOR ENABLING CAPABILITIES The opinions or assertions contained herein are the private views of the author and are not to be construed as official 26MAR2020 v.2</p> </div>

APPENDIX D: SLEEP CHECKLIST

<div> <div>COVID-19</div> <div>SLEEP CHECKLIST</div> <div>LEADERS AND SUPERVISORS</div> </div> <div>  </div>	
<p>Leaders and supervisors can take care of their staff and support team performance by prioritizing sleep for everyone.</p> <p>Use the SLEEP acronym...</p>	<ul style="list-style-type: none"> ➤ Set the conditions ➤ Lead by example ➤ Encourage sleep ➤ Educate about sleep ➤ Plan and prioritize
<p><u>SET THE CONDITIONS</u></p> <ul style="list-style-type: none"> <input type="checkbox"/> Ensure light boxes and black out curtains are available for staff <input type="checkbox"/> Designate appropriate, comfortable spaces for staff napping <p>Ask yourself: "Have I created a culture that supports sleep?"</p>	
<p><u>LEAD BY EXAMPLE</u></p> <ul style="list-style-type: none"> <input type="checkbox"/> Watch your own sleep habits and make sure you are getting enough sleep <input type="checkbox"/> Model appropriate caffeine use and sleep health <input type="checkbox"/> Acknowledge the reality of sleep debt and fatigue during COVID-19 <input type="checkbox"/> Avoid sending texts or emails to staff during non-duty hours <p>Ask yourself: "Am I walking the walk?"</p>	
<p><u>ENCOURAGE</u></p> <ul style="list-style-type: none"> <input type="checkbox"/> Ask your staff about their sleep <input type="checkbox"/> Emphasize the importance of sleep <input type="checkbox"/> Allow and encourage staff to take naps when appropriate <input type="checkbox"/> Talk about the importance of sleep at all levels of leadership <p>Ask yourself: "Am I checking in with my team about their sleep?"</p>	
<p><u>EDUCATE</u></p> <ul style="list-style-type: none"> <input type="checkbox"/> Reinforce points about self-awareness, caffeine, and light <input type="checkbox"/> Ensure team members know the basics of sleep health (e.g., 7-9 hrs per night) <input type="checkbox"/> Encourage staff to get sleep problems checked out medically as needed <input type="checkbox"/> Remember that decision-making and moral reasoning are impacted by lack of sleep <input type="checkbox"/> Remind your team that good sleep helps to protect health and fend off infection <p>Ask yourself: "What information am I sharing about sleep?"</p>	
<p>RESEARCH TRANSITION OFFICE, CENTER FOR ENABLING CAPABILITIES AND BEHAVIORAL BIOLOGY BRANCH, CENTER FOR MILITARY PSYCHIATRY AND NEUROSCIENCE, WALTER REED ARMY INSTITUTE OF RESEARCH • 06APR20 V.1 THE OPINIONS OR ASSERTIONS CONTAINED HERE IN ARE THE PRIVATE VIEWS OF THE AUTHORS AND ARE NOT TO BE CONSTRUED AS OFFICIAL.</p>	

RESEARCH TRANSITION OFFICE, CENTER FOR ENABLING CAPABILITIES AND BEHAVIORAL BIOLOGY BRANCH,
 CENTER FOR MILITARY PSYCHIATRY AND NEUROSCIENCE, WALTER REED ARMY INSTITUTE OF RESEARCH • 06APR20 V.I
 THE OPINIONS OR ASSERTIONS CONTAINED HERE IN ARE THE PRIVATE VIEWS OF THE AUTHORS AND ARE NOT TO BE CONSTRUED AS OFFICIAL.



NIGHT SHIFT TIP: LEVERAGE ANCHOR SLEEP

- Providers who cover night shifts may want to adjust their sleep to nighttime sleeping on their days off, but this change may cause havoc with their restorative sleep
- Instead, maintain “anchor sleep”: On their days off, make sure that at least 4 hours of sleep are anchored—or match—to their typical nightshift sleeping schedule
- However, it is best to keep providers on the same schedule as long as possible, as it is difficult for them to switch from day shifts to night shifts frequently

PLAN AND PRIORITIZE: SCHEDULING SHIFTS

- ☐ Limit staff shifts to 12 hours when possible
- ☐ Plan forward shift rotations that move with the clock
 - Shifting from day to evening, and evening to night, makes for an easier transition
- ☐ When possible, schedule shifts according to people’s chronotype
 - Put your “early birds” on the morning shift and your “night owls” on the night shift
- ☐ Don’t extend schedules for night shift workers
 - Have staff attend meetings and complete administrative tasks during their shifts
- ☐ Give team members more time to sleep after a long shift
 - The need for sleep goes up after longer periods of wakefulness

Ask yourself: “Am I scheduling my team members’ shifts effectively?”

PLAN AND PRIORITIZE: MANAGING SHIFT TRANSITIONS

- ☐ Stagger shifts by changing out some team members every 4 hours
 - This enables new team members to refresh the remaining team
- ☐ Ensure a team member who is shifting their schedule isn’t alone on the floor
 - Make sure others are around to keep them alert
- ☐ Prevent staff errors toward the end of a night shift
 - Establish additional safety protocols given the documented elevated risk in errors
- ☐ Allot time off for individuals who are significantly shifting their schedule
 - Allow a minimum of 32 hours off for those with an 8 hour change in shift time
- ☐ Check in with team members to see how their shift schedules are working for them

Ask yourself: “Are my team members handling shift changes safely?”

THIS IS A MARATHON, NOT A SPRINT.

BY PRACTICING SLEEP LEADERSHIP, YOU AND YOUR TEAM CAN ADVANCE THE MISSION TO COMBAT COVID-19.

Selected references: Barger et al. (2018). Effect of fatigue training on safety, fatigue, and sleep in emergency medical services personnel and other shift workers: a systematic review and meta-analysis. *Prehospital Emergency Care*, 22(sup1), 58-68. | Burgess et al. (2007). Optimal shift duration and sequence: recommended approach for short-term emergency response activations for public health and emergency management. *American Journal of Public Health*, 97(Supplement_1), S88-S92. | Harrison et al. (2019). Circadian Profile of an Emergency Medicine Department: Scheduling Practices and Their Effects on Sleep and Performance. *The Journal of emergency medicine*. | Patterson et al. (2018). Evidence-based guidelines for fatigue risk management in emergency medical services. *Prehospital emergency care*, 22(sup1), 89-101.

APPENDIX E: COVID-19 IN FORWARD-DEPLOYED SETTINGS EXECUTIVE SUMMARY

*Contributors (*Chapter Leads, †Main Editors)*

LTC Jason B. Alisangco, MC, USA*
 LT Emily A. Arnold, NC, USN*
 HM3 Maxwell C. Basden, USN
 HM2 James E. Bean, USN
 CDR Robert C. Call, MC, USN*†
 HM2 Tam V. Dang, USN
 LT Timothy D. Dye, NC, USN
 LT Amy L. Erdman, NC, USN
 HM1 Antoine S. Giacometti, USN
 LCDR Jason M. Hardwick, MC, USN*
 HM2 Victor Isarraraz III, USN
 CPT Rian N. Jockisch NC, USA
 LCDR Arthur T. Johnson IV, MC, USN

LT Scott E. Krasovetz, MSC, USN
 HM3 Sterling H. Magee, USN
 LCDR Andrew E. Mancusi-Ungaro, MC, USN*
 HM1 Christopher L. Moore, USN*
 LT Nicole M. Pendry, NC, USN*
 LCDR Kevin T. Petersen, MC, USN*†
 CAPT Craig J. Randall, MC, USN*
 HM2 Lucero E. Rocha, USN
 HM2 Kareem S. Rouse, USN
 CAPT Kristian E. Sanchack, MC, USN*
 LTJG Jose E. Santos-Calderon, NC, USN
 HMCS Johnny J. Stewart, USN*†
 HM2 Michael L. Ward, Jr., USN

EXECUTIVE SUMMARY

INTRODUCTION

From June to November 2020, a forward location experienced four discrete COVID-19 outbreaks. Each outbreak revealed new improvement points, response flaws, and potential COVID-19 exposures. Consequently, the COVID-19 response posture for the location matured as a result of the outbreaks. With over 400 COVID-19 positive cases and more than 1,500 completed quarantines, this Executive Summary seeks to share our observations, narrative, setbacks, and lessons learned managing COVID-19 in an operational, forward-deployed setting. We approached this paper with honesty and transparency, as many of our worst days became our greatest teachers. We also quickly realized that sustained eradication of COVID-19 on our base would require a level of operational lockdown incompatible with operational requirements. Consequently, our end-goal became defined by our ability to mitigate COVID-19 from compromising the joint-coalition mission and not the number of COVID-19 cases or time between outbreaks.

The following table and figures summarize the four COVID-19 outbreaks.

Table 1 - Summary of Four Forward Deployed Setting Outbreaks

Duration	Outbreak	# of Cases	Key Lessons Learned
10 Days	Single Camp	35	Communication, early symptom reporting, minimizing work area transmission
25 Days	Base-wide (from local national exposure)	210	Deliberate messaging, aggressive contact tracing and testing, COVID-19 source control, data-driven decisions and algorithms, team-based strategies
5 Days	Travel Quarantine	37	Small-cohort quarantine, out-of-quarantine testing
18 Days	Multi-Cluster Exposure	85	Targeted base amenity closures, efficacy of unit-level mitigation measures, intra-theater spread

Figure 2. Cumulative COVID-19 positive cases. The four distinct outbreaks are outlined in red





Figure 3. Forward Deployed Setting Daily COVID-19 case count. The four distinct outbreaks are outlined in red



The subsequent chapters are deliberately detailed and intended for the reader seeking to develop and implement the described measures and strategies. A more concise summary is provided through this Executive Summary and the bulleted lessons learned at the beginning of each chapter.

The chapters are broken into four sections. The lessons learned from each chapter are included here in the Executive Summary. Further details can be found in the respective chapters. The first two chapters discuss the background and setting of COVID in the forward deployed setting. Chapters 3-7 focus on the fundamental principles of COVID-19 risk mitigation. The next two chapters, chapters 8-9, highlight events and lessons learned from our experiences building and maintaining isolation and quarantine facilities.

The final two chapters discuss austere medical and nursing COVID-19 care, as well as MEDEVAC considerations. Finally, the appendices are for supplemental usage and include the lessons learned report by unit surgeon from the initial, single-camp COVID-19 outbreak.

FUNDAMENTAL PRINCIPLES OF COVID-19 RISK MITIGATION

Bluntly stated, our location was not adequately prepared for the COVID-19 outbreaks that began in June and escalated into late August and early September. While the initial, single-camp outbreak provided early observations and lessons learned, it may have contributed to a false sense of security that COVID-19 could be easily contained and controlled. The base-wide outbreak occurring in August quickly dispelled this belief. During the 25-day outbreak, our leadership discovered that it required a deliberate, multi-faceted strategy to prevent widespread infection. Specifically, we developed a COVID-19 risk mitigation plan built upon the following principles:

- Aggressive Contact Tracing and Testing
- Deliberate Messaging
- COVID-19 Source Control
- Data-Driven Algorithms and Decisions
- Team-Based Solutions

A comparison between the base-wide outbreak and the subsequent multi-cluster outbreak suggests the validity of our methods. The local national base-wide outbreak began with a single exposure point and spread across the base. Over 25 days, there were 210 positive cases from 23 different units. In contrast, the operational exposure outbreak had five independent exposure sources, yet only lasted 18 days and had 85 positive cases. Despite four additional “patient zeros,” the outbreak lasted seven days less with 60% fewer cases. Additionally, base amenity closures were significantly

reduced during the multi-cluster outbreak. Lastly, two units with a minimalist COVID-19 mitigation posture proved responsible for 51 of the 85 positive cases. With stricter internal mitigation practices in these two units, it is estimated that 60% of the cases could have been avoided.

AGGRESSIVE CONTACT TRACING AND TESTING LESSONS LEARNED

1. If cohort quarantine is undertaken, it is imperative to test all close contacts regardless of symptoms before group quarantine. A single asymptomatic COVID-19+ individual can spread the virus across an entire cohort. It also establishes the boundaries of the COVID-19 spread.
2. An analogy of wildfire containment is useful in explaining contact tracing objectives and policies. Wildfires are first contained and then extinguished.
3. Any personnel can be trained in contact tracing utilizing free online courses and forms.
4. Contact tracing is a joint responsibility between the medical team and the affected unit/company. Our best results occur when medical staff engage the unit/company leadership and jointly produce a close contact list.
5. Most close contacts can be identified with the following questions: Who are your roommates? Whom do you eat with? Whom do you exercise with? Whom do you work closely with? Can you think of any other persons with whom you were within 6 feet for a total of 15 minutes?
6. Utilization of mobile contact tracing teams for educating units, testing, evaluating high-risk work areas, or handling outbreaks in transient berthing where no unit chain-of-command exists is essential.
7. It is crucial to identify a team leader when coordinating contact tracing, testing, placement of individuals in appropriate quarantine or isolation facilities, and meals.
8. A centralized medical tracker is essential for maintaining accountability and documentation.

DELIBERATE MESSAGING LESSONS LEARNED

1. Patient education failed to achieve a wide-reaching distribution of knowledge as individuals immediately entered isolation or quarantine and were later reluctant to talk about their medical conditions.
2. Overhead base announcements proved an effective means of communication. Similarly, signs with posted COVID numbers outside the DFAC helped keep the public engaged, as did a base-wide catchphrase: "Don't let anybody close contact you."
3. A wide-distribution daily email with current case numbers, new case locations, narrative, and a daily COVID-19 question enabled general populace education, base situational awareness, and unified messaging.
4. A pamphlet addressing the long-term health risks of COVID-19 helped bridge the lack of information many patients had about COVID-19.
5. The use of a phone messenger application allowed real-time communications within the Role 2 and between the base medical assets, allowing for facilitated decision making, unified policies, and streamlined treatment.
6. Commanders, both local and theater-level, were able to make decisions based on up-to-date information. Daily counts of infections, quarantined folks, isolation folks, and PUIs enabled resources to match needs.
7. Collaboration with individual unit leaders was paramount in enforcing the strict but necessary guidelines and restrictions. The ultimate responsibility rested with unit leadership to ensure their personnel complied with required mitigation policies.

COVID-19 SOURCE CONTROL LESSONS LEARNED

1. Monitor external sources of data for trends of COVID-19 in the surrounding cities and countries.
2. You can learn more from close contact tracing by physically assessing the location as well as interviewing the exposed personnel.
3. Strict controls of movement on and off the base in conjunction with testing helped identify the local national sub-contractor population as a reservoir for COVID-19.
4. The basics rules of masking, avoiding large groups of people, and social distancing work well to minimize infectious spread but require constant messaging and monitoring due to human factors.
5. Frequent communications involving as many leaders as possible helped disseminate information. Public announcements and signage helped ensure the message was also pushed to the general base population.
6. Running the lab specimens posed a significant risk to lab personnel. Ensuring the lab spaces are safe, secure, and have adequate ventilation as well as a protective hood may decrease the risk of transmission.
7. Use of N95s for mission-critical workers successfully ensured the continuation of critical operations.
8. Decreasing the use of tents and increasing the use of containerized housing units (CHUs) will decrease the risk of exposure and help develop “neighborhoods” where units can work near where they live.
9. Quarantine tents should ideally be spaced appropriately and contain ten or fewer people per tent.

DATA-DRIVEN ALGORITHMS AND DECISION LESSONS LEARNED

1. Understand the limitations of the COVID-19-PCR test (false-negative rates).
2. Develop a testing strategy for when testing supplies are limited vs. plentiful.
3. Procure individual rooms for quarantined close contacts whenever possible.
4. Create a rolling average for cases per day, and establish a cutoff value to trigger base-wide action.
5. A “one-size fits all” base posture response is ineffective as every outbreak has unique dynamics, threat profile, and containment strategy.
6. Create a panel of medical experts on base who demonstrate an evidence-based approach to decision-making and understand epidemiology.

TEAM-BUILDING AND TEAM-BASED SOLUTIONS LESSONS LEARNED

1. Establish responsibilities of all base stakeholders: base leadership, medical leadership, medical staff, embedded unit medical personnel, and BOS-I.
2. Ensure broad, effective lines of communications among stakeholders that allow communication 24/7.
3. Identify POCs for every unit or contractor company. Medical personnel are preferred, but supervisors or leaders can substitute if no medical assets are attached.
4. Conduct frequent, established meetings with key medical staff (ISOFAAC mayors, mobile team leaders, lab, supply, etc.)
5. Train each unit on contact tracing, proper quarantine procedures, and support for personnel in quarantine or isolation.

ISOLATION AND QUARANTINE

Perhaps the most significant challenge with COVID-19 management was the implementation of CDC-guided isolation and quarantine practices. At its peak, our location had nearly 100 personnel in isolation and over 500 personnel quarantined. During the initial, single-camp outbreak, a CHU was converted into a COVID-19 isolation facility (ISOFAC) with 29 double-occupancy rooms. The CHU ISOFAC proved incapable of housing all the isolation patients, and an additional ISOFAC was established. Each isolation camp required medical oversight, logistic supplies, food delivery, and facility maintenance. Due to COVID-19 concerns, contracting support for these facilities was limited.

At the time of our arrival, the four open-bay quarantine tents comprising our quarantine capabilities were being dismantled due to a construction project. As of early July, we did not have a designated quarantine area. In just over a week, the base quarantine camp went from a “napkin sketch” to a functional quarantine camp with seven open-bay tents and 32 initial occupants. There were many setbacks and lessons learned, but the final collaboration between the medical assets, BDOC, BOS-I, Force Protection, and the base leadership proved a successful team in overcoming the ever-changing challenges.

ISOLATION FACILITIES LESSONS LEARNED

1. While both COVID-19 positive and PUIs require isolation, it is important not to mix them to prevent the spread of COVID-19 to a non-infected PUI.
2. 24/7 Communication between COVID-19 Isolation berthing and the role of care is mandatory as patients may develop worsening symptoms.
3. Design ISOFAC policies and areas to allow maximum autonomy and self-management by the ISOFAC occupants and their units, including cleaning, food delivery, and transport.
4. A designated “dead drop” area allows unit representatives to deliver food and supplies while avoiding inadvertent interactions with COVID positive patients.
5. Identify a camp mayor for each isolation facility. Camp mayors should become familiar with base policies and instructions for requested services, repairs, and maintenance.
6. Isolation facilities must include plans for medical or security evacuations including a designated shelter area.

QUARANTINE FACILITIES LESSONS LEARNED

1. Quarantine is a unit responsibility, though medical has much to offer in advisement and assistance for quarantine planning, tracking, and execution.
2. Establishing a base quarantine camp required strong partnerships between medical, BOS-I, BDOC, and base leadership.
3. Designating one or two days per week to start travel quarantine cohorts helps reduce limited quarantine resources and streamlines throughput.
4. Base camps require daily walkthroughs from a Camp Mayor to mitigate current or future concerns.
5. Daily vital sign checks for quarantined personnel increases the risk of close contact exposure with little benefit.

MEDICAL AND NURSING COVID-19 CARE

While most COVID-19 patients never required inpatient care, not all patients were so fortunate. During the first outbreak, 9 of the 35 COVID-19 patients were repatriated to LRMC. Twelve of the remaining patients required a 6-week follow-up for persistent COVID-19 symptoms, primarily fatigue and exercise intolerance. One of the twelve was found to have reduced cardiac ejection on point-of-care ultrasound testing and was MEDEVAC'ed for a full evaluation. During the base-wide outbreak, there were 10 COVID-19 related MEDEVACs and 16 cases of post-COVID-19 pericarditis (inflammation of the heart lining)

Every military base has a different medical footprint. Practical implementation of the lessons learned will depend on site and capability. Below, we offer our observations on providing medical, nursing, MEDEVAC, and ancillary care for patients who develop more than mild COVID-19 symptomology.

COVID-19 MEDICAL CARE LESSONS LEARNED

1. Patients with known or suspected COVID-19 symptoms should be kept separate from other patients and medical staff throughout their medical care, to include triage, resuscitation, and inpatient stay.
2. A standalone five-bed COVID ward/ICU and a separate COVID resuscitation facility allows COVID-19 patients to receive the same level of care as they would in the primary facility while protecting other patients, medical providers, and staff.
3. COVID-19 laboratory facilities should either include a hood for specimen processing and testing or be located in an independent lab annex.
4. Inadequate test kit supplies frequently threatened our response capability.
5. Outside X-rays may help prevent COVID transmission, though accommodations need to be considered for inclement weather.
6. DoD medication lists were beneficial in helping pharmacy stock appropriate medications.
7. With most COVID-19 cases demonstrating only mild symptoms, the OTC symptomatic medications proved the highest demand items.
8. Common OTC "go-bags" were supplied to patients in ISOFACs, preventing the need for a medical visit whenever an individual requested symptomatic treatment.

INPATIENT NURSING LESSONS LEARNED

1. Medical personnel should obtain proper fit testing with multiple styles N95 masks before deployment.
2. Ensure providers have access to properly fitted masks per fit testing results.
3. Separate COVID-19 facilities from normal daily operations.
4. Consider keeping operational areas "clean" by swabbing patients outside. If not possible, at least separate from the main facility to decrease exposure in close quarters.
5. Ensure measures to maintain and allow for social distancing, such as minimizing unnecessary staff at the patient care facilities.
6. Keep N95s out of the direct sunlight and away from the outdoor elements. Conserve PPE in a dry, temperature-controlled environment.
7. Create a bleach/water ratio for cleaning when Cavicide wipes are not available.
8. Do not use chairs or equipment that are fabric and cannot be appropriately sanitized.
9. Build your response on pre-established DoD, command, and CDC guidelines.
10. Recognize that with limited medical resources, it is important to wisely allocate time and personnel.