

## 13. Prophylaxis

- 13.1. \_\_\_\_\_ Protonix 40 mg IV Q day
- 13.2. \_\_\_\_\_ Lovenox 30 mg SQ BID OR \_\_\_\_\_ Heparin 5000 U SQ TID starting \_\_\_\_\_
- 13.3. \_\_\_\_\_ Pneumatic compression boots

## 14. Ventilator Settings

- 14.1. Mode: \_\_\_\_\_ SIMV \_\_\_\_\_ CMV \_\_\_\_\_ AC \_\_\_\_\_ CPAP
- 14.2. FiO<sub>2</sub>: \_\_\_\_\_ %
- 14.3. Rate: \_\_\_\_\_
- 14.4. Tidal Volume: \_\_\_\_\_ cc
- 14.5. PEEP: \_\_\_\_\_
- 14.6. Pressure Support: \_\_\_\_\_
- 14.7. Insp Pressure: \_\_\_\_\_
- 14.8. I/E Ratio: \_\_\_\_\_
- 14.9. \_\_\_\_\_ APRV: Phi \_\_\_\_\_ Plow \_\_\_\_\_ Thi \_\_\_\_\_ Tlow \_\_\_\_\_ FiO<sub>2</sub>: \_\_\_\_\_ %
- 14.10. \_\_\_\_\_ Maintain patient in soft restraints while on ventilator
- 14.11. \_\_\_\_\_ Wean FiO<sub>2</sub> to keep SpO<sub>2</sub> > 90-96% or PaO<sub>2</sub> 60-100 mmHg
- 14.12. \_\_\_\_\_ nebulizer/MDIs: \_\_\_\_\_ Albuterol \_\_\_\_\_ Atrovent \_\_\_\_\_ Xopenex Unit Dose Q 4 hrs

## 15. Analgesia/Sedation/PRN Medications

- 15.1. Analgesia/sedation goal is Richmond Agitation Sedation Scale (RASS), scale below, of 0 (alert and calm) to -3 (moderate sedation). Hold continuous infusion for RASS of -4 (deep sedation) or higher.
- 15.2. \_\_\_\_\_ Propofol gtt at \_\_\_\_\_ mcg/kg/min, titrate up to 50 mcg/kg/min.
- 15.3. \_\_\_\_\_ Fentanyl gtt at \_\_\_\_\_ mcg/hr titrate up to 250 mcg/hr; for analgesia may give 25-100 mcg IVP Q 15 minutes for acute pain or burn wound care.
- 15.4. \_\_\_\_\_ Morphine gtt at \_\_\_\_\_ mg/hr, titrate up to 10 mg/hr, for analgesia may give 2-10 mg IVP Q 15 minutes for pain or burn wound care.
- 15.5. \_\_\_\_\_ Versed gtt at \_\_\_\_\_ mg/hr, titrate up to 10 mg/hr ; may give 2-5 mg IVP Q 15 minutes for acute agitation or burn wound care.
- 15.6. \_\_\_\_\_ Ativan gtt at \_\_\_\_\_ mg/hr, titrate up to 10 mg/hr; may give 1-4 mg IVP Q 2-4 hours for acute agitation.
- 15.7. Important: Hold continuous IV analgesia/sedation at 0600 hrs for a RASS of -4 or -5. If further analgesia/sedation is indicated, start medications at ½ of previous dose and titrate for target RASS.
- 15.8. \_\_\_\_\_ Morphine 1-5 mg IV Q 15 minutes prn pain
- 15.9. \_\_\_\_\_ Fentanyl 25-100 mcg IV Q 15 minutes prn pain
- 15.10. \_\_\_\_\_ Ativan 1-5 mg IV Q 2-4 hrs prn agitation
- 15.11. \_\_\_\_\_ Percocet 1-2 tablets po Q 4 hrs prn pain
- 15.12. \_\_\_\_\_ Tylenol \_\_\_\_\_ mg / Gm PO / NGT / PR Q \_\_\_\_\_ hrs PRN for fever or pain
- 15.13. \_\_\_\_\_ Morphine PCA; Program (circle one): 1   2   3   4
- 15.14. \_\_\_\_\_ Zofran 4-8 mg IVP Q 4 hrs PRN for nausea/vomiting
- 15.15. \_\_\_\_\_ Dulcolax 5 mg PO / PR Q day PRN for constipation

## 16. Specific Burn Wound Care

- 16.1. Cleanse and debride facial burn wounds with Sterile Water or (0.9% NaCl) Normal Saline Q 12 hrs, use a washcloth or 4x4s to remove drainage/eschar
- 16.2. Cleanse and debride trunk and extremities with chlorhexidine gluconate 4% solution (Hibiclens) and Sterile Water or Normal Saline, before prescribed dressing changes
- 16.3. Change fasciotomy dressings and outer gauze dressings daily and as needed; moisten with sterile water Q 6 hours and as needed to keep damp, not soaking wet.

## 17. Other Orders

- 17.1. \_\_\_\_\_
- 17.2. \_\_\_\_\_

- 18. **Notify Physician if:** SBP < \_\_\_\_\_, MAP < \_\_\_\_\_, HR < \_\_\_\_\_ or > \_\_\_\_\_, SaO<sub>2</sub> < \_\_\_\_\_%, T > \_\_\_\_\_, UOP < 30 mL/hour for 2 consecutive hours